

# Sorghum as a Model Crop for Drought Stress Tolerance

**Mulatu Gidi**

Ethiopian Institute of Agricultural Research, National Agricultural Biotechnology Research Centre, Holeta, Ethiopia

**Email address:**

mulatugi23@gmail.com

**To cite this article:**

Mulatu Gidi. Sorghum as a Model Crop for Drought Stress Tolerance. *Advances in Bioscience and Bioengineering*. Vol. 11, No. 3, 2023, pp. 54-65. doi: 10.11648/j.abb.20231103.14

**Received:** August 4, 2023; **Accepted:** August 23, 2023; **Published:** September 8, 2023

---

**Abstract:** Sorghum is one of the most significant C4 cereal crops grown in dry and semi-arid regions of the world. It is a major staple crop for millions of people in Sub-Saharan Africa and South Asia. Drought is an important constraint on agricultural production and productivity around the world. It has a significant impact on plant growth, development, and yields. Drought stress risks food security by having a substantial impact on sorghum growth and development, grain yields, and nutritional quality. Sorghum has become known as a drought-tolerant model crop when compared with many other crops. Its ability to withstand extreme environmental conditions makes it a feasible model crop for studying abiotic stress responses and developing stress-tolerant crops. Sorghum response and/or tolerance mechanisms include morphological, physiological, and molecular changes. Drought stress tolerance mechanisms in sorghum include drought escape, early flowering, stay-green, drought avoidance, leaf area, osmotic adjustment, stomata-mediated drought responses, cuticular wax production, root characteristics, and drought tolerance. Biotechnology and its advanced approaches, such as QTL, marker-assisted backcrossing, genetic engineering, and others, are used for screening drought-tolerant genotypes that can withstand drought stress. Therefore, focusing on the drought-tolerant genotypes will boost the speed of the sorghum breeding program, which will feed millions of people worldwide, particularly in Sub-Saharan Africa.

**Keywords:** Drought, Drought Tolerance, Grain Yield, Sorghum, Stay-Green

---


## 1. Introduction

Sorghum (*Sorghum bicolor* L.) is an annual grass in the Poaceae family with a C4 metabolism. The crop originated between 8,000 and 5,000 years ago in Northeastern Africa [1]. It is a photosynthetically proficient C4 plant that provides enormous amounts of grains, forage, sugars that ferment, and cellulosic fibers for a number of uses that encompass bioenergy and bio-industrial feedstocks [2]. After maize, rice, wheat, and barley, sorghum is the world's fifth major cereal crop [3]. In 2020, global sorghum production was 29.8 million metric tons (MMT), with a production rate of 1.5 metric tons per hectare (t ha<sup>-1</sup>) [3]. Global sorghum production was estimated at 60.06 million metric tons in 2021–2022. Nigeria stands first in total production with 7 million metric tons (12%), followed by the US, Sudan, and Mexico (Figure 1). It is a food security crop for about 500 million people in nations with low incomes, mostly in dry and drought-prone areas where moisture stress is significant. [4].

Drought is a recurring incident that threatens agricultural productivity and has a global impact on the livelihoods of people [5]. It has an impact across many African sorghum-producing countries [6]. Drought is anticipated to become more prevalent in the future due to climate change [7]. Nevertheless, the consequences of drought on agricultural output depend on rainfall distributional trends compared to overall annual rainfall. As a result, drought is nowadays a serious issue for crop development and growth, particularly in tropical countries [8]. Drought stress impacts nearly all stages of plant development; nevertheless, the emergence of seeds and the early seedling growth phase [9] as well as reproduction phases, particularly in sorghum [10], are extremely vulnerable and crucial. It reduces plant health, development, and growth and reduces production by limiting carbon absorption, stomatal conductance, and cell turgor [10, 11]. The wilting of leaves and loss in leaf surface indicate moisture stress indications on crop plants, as are bud or flower production, sink number, total development, and crop yield [10, 12]. Understanding the impact of drought on crops is therefore critical for generating better varieties with

consistent high crop production. Crop responses to drought stress, on the other hand, are multifaceted and differ depending on climatic circumstances, the frequency and

length of the stress, the plant's species and diversity, and the plant's physiologic stages at the time of the stress occurred [13].



**Figure 1.** Global sorghum production from 2021–2022 (Source: USDA, [fas.usda.gov](https://fas.usda.gov)).

## 2. Sorghum as a Model Crop

Sorghum bicolor, a C4 cereal crop used for food, feed, and biofuel, has been found to be drought tolerant [14, 15]. It is a major source of food in developing countries, several of which are particularly vulnerable to climate change [16]. Sorghum was cultivated in Africa 8000 years ago, and its adaptation to arid environments shows that drought tolerance developed at domestication [16–18]. Sorghum is a drought-resilient crop due to its domestication in dry environments. Therefore, it is an ideal model crop for studying drought-stress mechanisms of response in an agriculturally and economically valuable cereal crop. Furthermore, the variation in genetic makeup within sorghum genotypes or accessions, as well as its relatively small diploid genome and close relationship to *Zea mays*, make it a good candidate for identifying and clarifying drought tolerance genetic processes in related grasses [19–21].

## 3. Importance of Sorghum

Sorghum is an essential food source in many dry, semi-desert, and tropical areas around the globe due to its being more tolerant and prolific than other crops when exposed to diverse environmental conditions. It is abundant in fiber and protein and also contains more calcium, phosphorus [22], and potassium than rice or wheat [23]. Sorghum possesses essential human health and nutritional properties, and the fact that it is gluten-free makes it an ideal choice as an option to wheat for use in the making of special meals for people with celiac disease who are gluten-sensitive [24, 25]. It consists of resistant starch, which may decrease digestion, particularly in orphan, but is significant in the

fight against human obesity and as a source of food for diabetics [26]. Sorghum also contains a lot of fiber and bioactive components such as tannins, phenolic acids, anthocyanins, phytosterols, and policosanols. These phytochemicals are desired in human food due to their antioxidant activity, which benefits health. Certain sorghums contain a high concentration of unique phytochemicals that have anti-cancer effects as well as producing bran with more antioxidant properties than fruits or berries [24, 25, 27, 28]. Sorghum is a key grain crop that is also used to make alcohol, bioethanol, and fuel, not just food.

## 4. Drought Stress

Drought remains one of the most important abiotic factors affecting crop production globally [29]. It reduces plant productivity by causing the closure of stomatal pores and lowering the photosynthesis process as well as growth [30]. Plant adaptability and reaction to drought are consequences of complex biological mechanisms such as physiological, biochemical, genomic, proteomic, and metabolomic alterations [31]. Drought-sensitive sorghum cultivars showed stress-related physiological alterations, including a shift in photosynthetic rate [32]. According to the research [33], drought stress lowers stomatal conductivity and evaporation rates, raises the quantity produced and leaves the temperature, reduces chlorophyll and Rubisco, increases O<sub>2</sub> evolution, and decreases PEPCase function in sorghum. According to multiple studies [32, 34], drought-tolerant sorghum cultivars had significantly higher Fv/Fm and photosynthesis rate values. Besides their capacity to escape or withstand drought, photosynthesis restoration on rehydrate is critical in determining crop drought tolerance and avoiding grain yield

loss [35]. Transpiration efficacy in drought-resilient sorghum cultivars was similar across normal and drought-stressed crops; however, in drought-susceptible genotypes, there was a substantial statistical difference between control and drought-stressed plants [32]. Furthermore, during the stress caused by the drought period, cultivars that proved greater resilience to stress had a significantly greater WUE than those that were more sensitive [32]. In sorghum, grain yield has been observed to be significantly associated with water removal and absorption efficiency [36]. Drought-resistant cultivars maintain moisture for use during the grain-filling stage in drought-prone regions and have lower stomatal conductivity and transpiration levels at the vegetative stage [37].

## 5. Impacts of Drought Stress on the Sorghum Crop

Sorghum, as a significant crop for climate change resilience, is outstanding for smallholder farmers in Africa and Asia who have no access to irrigation to sustain food security [38]. Climate change has resulted in the instability of weather patterns, particularly in arid areas where the duration of dry periods has increased, resulting in crop wilting and food shortages. Figure 2 shows the impact of drought stress on sorghum plants in Ethiopia.



**Figure 2.** Impacts of drought stress on sorghum in Ethiopia, Photos—January 2016. (Source: Adebbe et al., 2017).

### 5.1. On Sorghum Growth and Development

Drought stress will have a considerable impact on plant growth early in the growing season. Seedling deaths are of great concern in arid regions where drought damage is prevalent, and they are most severe during seedling emergence and maintenance under the combination of extreme drought and high-temperature conditions [39]. Drought can cause standing mortality in the sorghum following entire emergence but before seedling establishment [40]. Drought stress is possibly affecting the initial stages of plant growth (germination, emergence, and seedling establishment). Consequently, the impact of drought-induced moisture deficits on the initial embryonic stages of sorghum

has attracted considerable attention. Nevertheless, there are significant differences in sorghum genotypes' reactions to varying degrees of drought-related stress. Drought stress can be detrimental to plant development at the beginning of the growth period; however, when late rainfall rates are sufficient, plants recover rapidly [41]. It reduces the seedling's vigor, germination rate index, and percent germination by increasing the respiration rate, which affects starch synthesis and energy generation (adenosine triphosphate (ATP)) [40]. According to the research [42], when drought stress was subjected, the shoot and root lengths of drought-sensitive genotypes were shorter than those of drought-tolerant genotypes.

### 5.2. On Sorghum Grain Yield

Drought stress affects grain yields through decreasing in seed size, quantity, or weight per panicle, as well as other agricultural characteristics [43]. Furthermore, it raises both direct and indirect agricultural expenses, decreasing yield and productivity. Despite the reality that sorghum is one of the most resilient drought-tolerant plants, adapted to a wide range of agroecology as well as low-input food production, extreme drought might result in considerable yield losses [44]. Grain yields in water-stressed locations are often reduced due to unexpected and insufficient rainfall [45]. According to the research [46], in a sorghum study, drought stress affects grain quality and yield both before and after flowering. Drought stress during the pollination stage could result in a considerable drop in the yield of grains due to the failure of egg implantation within the ovaries [43]. It may also limit the quantity of grain per panicle, which has a direct effect on grain yield during the flowering stage [47]. Drought stress during the post-flowering period lowered grain yield by roughly 50%, according to a two-year classic study involving 30 sorghum varieties [48].

### 5.3. On Sorghum Nutritional Quality


Several studies have shown how drought stress affects the nutritional content and composition of sorghum, resulting in crop alterations. It alters the interactions between morphophysiological traits on the one hand and source action and sinking strength on the other [49], alters grain physical and chemical properties [50], and decreases nutrients, mineralization, and membrane permeability [51]. Growing sorghum genotypes in dry environments, for example, decreased grain micronutrient content (Zn, Fe, Mn, and Cu) [50]. Drought stress resulting during the phase of flowering caused lower total starch, amylase, and amylopectin accumulation, which corresponds to the activity of enzymes on sugar nucleotide precursors during the filling of grains, such as starch synthase (SSS), granule-bound starch synthase (GBSS), starch branching enzyme (SBE), and starch debranching enzymes (DBE). In other studies, it increased the amount of total protein while decreasing the total soluble carbohydrate, crude protein, and proline concentrations in sorghum [43].



## 6. Drought Stress Tolerance Mechanisms in Sorghum

Plants have a variety of morphological, physiological, and biochemical adaptation processes in response to drought in order to thrive and live in harsh environments. Drought

escape, drought avoidance, osmotic adaptations, staying green, leaf area, waxiness on the stem, root shape and architecture, transpiration efficiency, and solubility solute secretion are all significant mechanisms in drought tolerance [52].



**Figure 3.** Diagrammatic depictions of the morphological, physiological, biochemical and molecular responses of sorghum to drought stress. This figure was created with BioRender (<https://biorender.com/>).

### 6.1. Drought Escape

Drought escape is a typical adaptation strategy that enables a crop to end its entire lifespan prior to a drought happening. Drought-resistant varieties end their lives prior to the appearance of extreme drought stress. During several field trials, an early flowering phenotype is used as a primary drought escape selection index for yield under drought conditions. A drought-escape sorghum phenotype is a germplasm that exhibits distinct responses to photoperiodism and homeostasis for bearing time [53].

### 6.2. Drought Avoidance

Drought-affected crops prevent dehydration by retaining significantly greater tissue levels of water, whether through increased intake of water or reduced loss of moisture through transpiration. Plants or cultivars with an earlier phenotype are known to be referred to as water spenders, whereas those with the latter are referred to as water savers. Water-spending plants survive droughts by expanding their root length, which assists in water absorption. Water savers, on the other hand, cover a portion of their stomata in order to reduce transpiration water loss and save soil moisture. Furthermore, drought stress is avoided by activating the osmotic adjustment mechanism, which includes solute buildup [10,

54].

### 6.3. Early Flowering

Drought stress during flowering decreases silk elongation, lengthens the anthesis silking delay, and reduces ovule fertilization and kernel abortion. Early flowering and short reproductive periods are essential characteristics under extreme drought circumstances because they reduce exposure to dehydration during the critical flowering and post-anthesis grain-filling periods [55]. In environmental settings with insecure rainfall or terminal dryness, flowering before the start of a development restricting drought is a key approach to guaranteeing yield production.



**Figure 4.** Early-flowering sorghum varieties (Source: online, March 16, 2016 From a Press Release)

#### 6.4. Leave Area

Drought stress causes varying responses in water consumption efficiency in leaves. Water usage efficiency at the level of the leaf is intimately tied to the physiological mechanisms controlling CO<sub>2</sub> and H<sub>2</sub>O gradients between the leaf and the air around the leaf [56]. Optimal leaf area is critical for optimum photosynthetic performance in a drought scenario. Furthermore, in drought-prone areas, optimum leaf area is critical to achieving high dry matter content and grain yields [57]. Sorghum's drought resistance was greatly influenced by leaf area. In regions prone to drought, the leaf area of sorghum is reduced to reduce water loss through evapotranspiration. Plant consumption of water and biomass buildup must take into account soil water evaporation rate, transpiration from the leaves, and plant growth style. Implementing techniques that minimize soil water evaporation and shift more water towards transpiration can improve water usage efficiency at the canopy level [56].



**Figure 5.** Photograph showing the stay-green trait in sorghum the field in Petancheru, India. The photograph was taken at the post-flowing stage with no supplemental irrigation. Sorghum plants on the right have the stay-green trait (B35), whereas plants on the left are senescent varieties (R16) (Kumar et al., 2011).

#### 6.5. Stay Green

Plants expressing stay green to keep functioning leaves that are green for an extended period of time as water shortage happens in the filling of grains [5]. Stay green is a moisture-stress adaptability trait seen in sorghum and other crops. Stay-green sorghum genotypes can preserve grain filling in drought condition. Grain yield and stay-green are goal qualities in grain sorghum breeding for wide adaptability to varied environments [58]. Staying green has been shown to improve tolerance of post-flowering dryness in sorghum grain. A number of QTLs linked with stay green have been found in sorghum; consequently, improving transmission of this trait into appropriate genetic backgrounds is critical [59]. According to the research [60], pre-flowering canopy variables were slightly connected with stay-green values for leaf senescence, while canopy factors had a considerably larger linkage with leaf senescence. This finding revealed that canopy size before flowering had little

effect on the production of a stay-green phenotype after flowering. The examined cultivars' resistance strategies included accelerated leaf senescence to reduce water loss and increased root length density to explore the deeper soil layers for moisture [60]. Sorghum line B35, a descendant of a hybrid between Ethiopian durra and Nigerian landrace, is a highly potential line carrying the 'stay green' phenotype and is often used as a trait source across regions [61]. Thus, green sorghum helps increase grain yields in both well-irrigated and drought-prone environments.

#### 6.6. Osmotic Adjustment

Osmotic adjustment is defined as a net solute increase that results in a decrease in tissue osmotic potential. Both osmolytes can accumulate in both the roots and the shoots, but their accumulation in both tissues serves different functions [39]. Water flow from higher concentration (soil) to lower concentration (tissue) happens as the water potential decreases due to decreasing osmotic potential [62, 63]. Osmotic adjustment was shown to be highly associated with deep root systems in sorghum, as indicated by the research [64, 65]. Sorghum leaves adjusted osmotically at a rate of at least 0.1 MPa per day as the lowest leaf water potential declined at a rate of around 0.15 MPa per day [66]. The shoots' osmotic adjustment prevents wilting and maintains relative water content and stomatal conductance [67, 68].

#### 6.7. Stomata-Mediated Drought Responses

Stomata serve a crucial role in plant carbon absorption as well as water loss processes. The cultivars physiologically adjusted to earlier water limits by closing their stomata, which decreased their transpiration and photosynthetic rate [60]. Stomata in drought-stressed plants are likely to close during midday vapor pressure deficit (VPD) and open primarily in the early morning when temperatures are lower before VPD becomes too high [69]. This function protects the plant's tissue hydration status, minimizing excessive loss of water through evaporation. Variations in leaf temperature can also be used to monitor this process. Despite stomatal closure avoiding transpirational water loss, it lowers CO<sub>2</sub> availability, impacting photosynthesis, plant development, and grain yield [70, 71]. Non-stomatal alterations affecting the rate of photosynthesis include reduced chlorophyll content, which decreases the plant's ability to collect sunlight and affects assimilation formation and transportation [21, 70].

#### 6.8. Cuticular Wax Production

Cuticular wax is the primary barrier that covers above-ground tissues in numerous species of plants, and it plays essential roles in plant abiotic and biotic stress tolerance. It is now implicated in defensive mechanisms against extreme UV radiation, high temperatures, bacterial and fungal diseases, insects, high salinity and low temperature [72]. Sorghum generates epicuticular wax (EW) in its leaves, sheaths, and culms. Epi-cuticular wax reduces transpirational and non-transpirational (nonstomatal) water loss and defends

plants from extreme drought stress, besides providing insect pest resistance [73]. A recent study demonstrated that short RNAs (sRNAs), such as microRNAs and small interfering RNAs, played essential roles in stress-induced plant adaptation. Furthermore, leaf cuticle and surface wax have crucial resilience to stress attributes, and several regulator genes regulate wax formation [74].

### 6.9. Root Traits

Plants are more vulnerable to harsh drought conditions, and roots perform key functions in plant survival, development, and reproduction. Plant roots have evolved to be the first organ that monitors and adjusts for shifts in soil moisture on morphological, anatomical, and molecular scales [75]. Roots serve as the vegetative organs in charge of water and nutrient intake as well as interaction with the soil microbes, and they play a vital role in dry tolerant to stress [76]. Roots have developed a capacity to modify their root structure architecture (RSA) characteristics when subjected to water stress [77-80]. Long and branched roots are one of the fundamental and indispensable traits for crops to tolerate droughts [81]. Small, fine root diameters, long, particular root lengths, and root length density at depths in soil with sufficient moisture are root traits associated with plant production during moisture stress [82]. As a result, roots play an important role in plant life because they take soil resources from deeper soil layers, and phenotyping allows for the comprehension of diversified root properties [83].

Sorghum gains access to more water via deeper root systems [5]. In dry circumstances, low xylem diameters in targeted seminal roots preserve water from the soil deep below the soil surface to be utilized for plant development, resulting in increased crop yields. Larger root growth angles result in root elongation towards deeper soil layers, which is regarded as an essential characteristic for drought access to and collection of deep soil water [84, 85]. Under drought-related condition, smaller diameter roots, higher SRL, and enhanced root hair density or length can enhance plant water uptake while reducing the plant carbon investment required for that uptake. Furthermore, plant hydraulic performance under drought stress could have been enhanced by greater capacity for nocturnal refilling of embolized xylem and alterations in inter-vessel pit structure to prevent turbulence [82]. Understanding root changes in response to water scarcity at many levels may thus offer a basis for choosing and growing drought-tolerant crop varieties.

### 6.10. Drought Tolerance

Sorghum is well-known for its capacity to withstand stress caused by drought, both intermittent and terminal. Tolerance behavior is influenced by the presence of a complicated and prominent root system, the ability to maintain an adequate amount of stomatal conductance over time, and the maintenance of internal tissue water potential via osmotic

adjustment and phenological flexibility [10, 54, 86].

## 7. Role of Biotechnology in Sorghum Tolerance to Drought Stress

Various biotechnology techniques have been used in sorghum crop improvement for drought tolerance. Marker-assisted backcrossing (MABC), quantitative trait loci (QTL), genetic engineering, and others are the most commonly used. Crop gene transformation techniques have been applied to identify and transfer the genes relevant to drought tolerance.

### 7.1. Marker-Assisted Backcrossing in Sorghum

Backcrossing is the transfer of alleles at one or more loci from the donor parent to the elite recurrent parent [87]. Backcrossing can be tedious and time consuming; the use of molecular markers has been important in identifying QTLs related to the expression of traits of interest. For example, several markers (simple sequence repeats and single nucleotide polymorphisms) have been chosen for mapping QTLs of stay-green in sorghum. These markers simplify breeding procedures by lowering the time it takes to release new, superior varieties and reducing breeding cycles. After six backcrosses, the introgression of elite trait alleles is achieved, with 99.2% genome recovery of recurrent parents [87]. Due to climate change in most regions of the world, researchers have focused on generating new drought-tolerant cultivars. A variety of molecular markers, such as delayed leaf senescence, leaf rolling, chlorophyll content, water-use efficiency, and yield, have been used to discover QTLs associated with stay-green in sorghum [88-90]. Some QTLs are found in all habitats, while others are found only in a few [91]. Backcrossing with an abundance of polymorphic, high-resolution, high-throughput, co-dominant, and informative molecular markers, such as SSRs and SNPs, is suggested to leverage QTLs from donor parents to recurrent parents [92]. SSR markers, for example, have identified Stg1-Stg4 QTLs that account for 10–30% of the phenotypic variation in drought tolerance in sorghum (Table 1). ICRISAT recently used SNP markers in sorghum to map Stg3A and Stg3B QTLs in chromosomes. These QTLs are very successful at delaying leaf senescence during post-flowering drought stress. In Tanzania, Mwamahenje, A. et al. inserted Stg3A and Stg3B QTLs into farmers' preferred sorghum varieties [93]. The QTLs were successfully introgressed by SNP markers snpSB00049, snpSB00053, and snpSB00054 for the Stg3A QTL and snpSB00101, snpSB00102, and snpSB00103 for the Stg3B QTL, which accounted for 30% of the grain yield increase and stay-green in water-stressed conditions. However, several of the markers failed to discover QTLs, indicating the need for more Stg3A and Stg3B mapping to diversify important drought-tolerance improvement.

**Table 1.** Molecular markers and stay-green QTLs for drought tolerance in sorghum.

Molecular markers	QTL	Position in Chromosome	PV (%)	Reference
Xtxp114, Xtxp38, xiabx378, SSR markers	Stg1	SB1-03	20	[94, 95]
XnhbsFCIIP67, Xtxp120, Xtxs584, and Xtxp31, SSR markers	Stg2	SB1-03	30	[94]
Xtxs1307, Xtxs1111, Xtxp1, Xtxp56, Xtxp286, SSRs markers	Stg3	SB1-02	16	[90]
Xtxs713, Xtxs387, Xtxp225, Xtxp15, SSR markers	Stg3	SB1-05	10	[90, 96]
snpSB00049, snpSB00053, and snpSB00054, SNPs markers	Stg3A	SB1-02	31	[93, 97]
snpSB00101, snpSB00102, and snpSB00103, SNPs markers	Stg3B	SB1-02	31	[93, 97]

## 7.2. QTL Pyramiding for Drought Tolerance in Sorghum

QTL pyramiding is the process of crossing one NIL to another, constructing distinct beneficial QTLs, and then using marker-assisted selection to develop novel lines with both helpful traits. Pyramiding QTLs for stress-tolerant traits and yields is advised with progressive increases in warmth and lack of moisture, particularly in arid environments, although the pyramiding approach is tedious and time-consuming [98]. Transfer of desirable traits from donor parents to the recipient is advised for the same species [99]. Drought tolerance in sorghum is improved by QTL pyramiding; nonetheless, efficiency in a well-irrigated condition is greater than performance in a water-stressed condition. Introgression lines with Stg3 and Stg1+2, for example, reduce yield by 10% in water-stressed conditions compared to well-irrigated conditions, showing better tolerance than other Stgs, which show 18–23% as moderate tolerance in sorghum [100]. Nevertheless, findings show that under drought stress conditions, crude protein, total soluble carbohydrate, and proline contents raise; this increase results in an improvement in grain quality nutrition, which enhances human health [43]. Pyramiding is the combination of QTLs with varying effectiveness to improve the expression of stay-green characteristics. The use of molecular markers assists in the detection of QTLs that may be pyramided to improve drought tolerance in sorghum [96]. Pyramiding of QTLs may fail to enhance sorghum stay-green due to incompatible gene activity, which prevents the expression of the traits needed. Pyramiding success is dependent on genetic architecture and accurate mapping of trait QTL [100]. Only the best expression of the trait of interest is chosen for further investigation in multi-location trials before being approved as a new variety during QTL pyramiding.

According to the research [101], locating QTLs for stay-green expression using molecular markers is critical for pyramiding to increase the heritability of new lines. Every QTL contributes a modest percentage to stay-green trait expression, which requires pyramiding to combine QTLs for expression. Pyramiding of stay-green QTLs in sorghum during pre-flowering and post-flowering drought increases the potential of producing new, promising lines with desirable traits through marker-assisted backcrossing [102, 103]. These findings help plant breeders assess the potential QTLs of introgressed traits that can be generated following pyramiding [104]. For successful pyramiding, appropriate molecular markers for mapping QTLs must be used to facilitate sorghum molecular breeding programs. Only

pyramided QTLs that show trait expression are suggested for selection in sorghum improvements in the pyramiding study.

## 7.3. Genetic Engineering for Drought Tolerance in Sorghum

The application of genetic engineering techniques holds great promise for enabling the generation of transgenic crops with little or no influence on the plants' physiological and morpho-biochemical characteristics. Most of the genes linked to drought tolerance are also linked to resilience to other environmental conditions [105]. Genetic transformation is an effective method of improving cultivated plant genetics. Tissue culture challenges, such as phenolic pigment accumulation and low regeneration frequency, have slowed progress in sorghum transformation. Longer durations of screening required for the recovery and regeneration of putative transgenic crops frequently hampered sorghum transformation condition optimization. The presence of methylation of the DNA in sorghum cells, which inhibits the functioning of transmitted genetic material, is likely responsible for sorghum's poor transferability [106]. Genetic engineering methods are particularly essential for modifying protein and starch levels, vitamin and micronutrient concentrations, protein fraction nutritional value, and resistance against environmental stresses [107].

## 7.4. Multi-Omics Research Advances in Drought Stress Tolerance in Sorghum

Water is one of the most basic needs for the development of plants. Plants adapt to water trouble by changing their morphology, physiology, and biochemistry in order to cope with direct and indirect consequences and live and/or flourish [108-110, 31]. Drought reactions in plants are categorized into three categories: drought escape, avoidance, and tolerance, which have been thoroughly investigated in previous studies [111, 112, 71]. To investigate and mine essential genes linked with strong drought tolerance in sorghum, RNA-seq was used to profile the up-and down-regulated genes linked with drought stress in the drought-tolerant sorghum cv. XGL-1, revealing that numerous genes with differential expression in the roots have been increased in functions such as sucrose metabolism and the raffinose family oligosaccharide biosynthetic process [113].

Functional investigations in sorghum indicated that the SbHKT1; 4 gene regulates cellular ion homeostasis and improves tolerance to drought and salt stress [114]. Various research efforts have recently used the comparative technique to discover DEGs across stress-tolerant and stress-sensitive

sorghum genotypes. A transcriptomic comparison of drought-sensitive and drought-tolerant genotypes (IS20351 and IS22330) revealed that the former responded to stress by hydrolyzing carbohydrates in roots, whereas the latter gained tolerance by increasing the synthesis of anti-osmotic agents and antioxidants (e. g., proline, betaine, and glutathione) [32]. Proteomics has also been applied to analyze the protein dynamics of sorghum roots in response to osmotic stress caused by PEG [115]. During drought stress, the concentrations of MDA and proline were steadily raised, as were the activities of superoxide dismutase (SOD), peroxidase (POD), and polyphenol oxidase (PPO). Proteomic analysis of osmotic-induced proteins in BTx623 roots detected proteins involved in protein synthesis, degradation, and defense [115]. These proteomic studies in sorghum are likely to show that cultivars that are drought-tolerant tend to up regulate their stress-related signaling, protein synthesis, and antioxidant activity to acquire improved drought tolerance.

## 8. Conclusion

Drought is a complex phenomenon that has an impact on agriculture by reducing its production all over the world. Plants respond to drought stress in a number of ways. Sorghum has higher resilience to unfavorable environmental conditions than other grain crops. Sorghum is a drought-tolerant grain crop that is used as food for millions of people living in dry-land areas. It can provide good yields even when water is scarce. Sorghum is regarded as an important crop species for studying the genetic and physiological mechanisms underlying drought tolerance in higher plants. Biotechnology, with its new and advancing approaches, can be used to overcome the problem of drought stress in crops. Due to the small genome size of sorghum, advanced biotechnology techniques have proven reliable for identifying genes related to more complex drought tolerance. Because of its potential to withstand drought under harsh climate change, it represents the future hope crop that will be the source of food for billions of people globally.

## Conflict of Interest

The author has not declared any conflict of interests.

## References

- [1] Mann, J. A., Kimber, C. T., and Miller, F. R. (1983). The origin and early cultivation of sorghums in Africa. Texas FARMER Collection.
- [2] Boatwright J. Lucas, Brenton Zachary W., Boyles Richard E., Sirjan Sapkota, Myers Matthew T., Jordan Kathleen E., Dale Savannah M., Nadia Shakoor, Cooper Elizabeth A., Morris Geoffrey P. and Stephen Kresovich, (2021). Genetic Characterization of a Sorghum bicolor Multiparent Mapping Population Emphasizing Carbon-Partitioning Dynamics. Genetics Society of America 1-14.
- [3] FAOSTAT, (2020). Food and agriculture organization of the United Nations. Rome, Lazio, Italy: FAO. Available at: <https://www.fao.org/faostat/en/#data/QCL>
- [4] Ejeta, G. (2005). "Integrating biotechnology, breeding, and agronomy in the control of the parasitic weed striga spp in sorghum," in The wake of the double helix: from the green revolution to the gene revolution, Bologna Bologna. Eds. R. Tuberosa, R. L. Phillips and M. Gale. (Bologna: Avenue Media) 239–251.
- [5] Liedtke J. D., C. H. Hunt, B. George-Jaeggli, K. Laws, J. Watson, A. B. Potgieter, A. Cruickshank and D. R. Jordan, (2020). High-Throughput Phenotyping of Dynamic Canopy Traits Associated with Stay-Green in Grain Sorghum. Plant Phenomics. <https://doi.org/10.34133/2020/4635153>
- [6] Eggen, M.; Ozdogan, M.; Zaitchick, B.; Ademe, D.; Foltz, J.; Simane, B. (2019). Vulnerability of sorghum production to extreme, subseasonal weather under climate change. Environ. Res. Lett. 14, 045005.
- [7] IPCC, Intergovernmental Panel on Climate Change (2013). Summary for policymakers. In Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to V Assessment Report of the Intergovernmental Panel on Climate Change; Stocker, T. F., Qin, G. K., Plattner, M., Tignor, S. K., Allen, J., Boschung, A., Nauels, A., Xia, Y., Bex, V., Midgley, P. M., Eds.; Cambridge University Press: Cambridge, UK, 2013.
- [8] Xu, Z.; Zhou, G.; Shimizu, H. (2018). Plant responses to drought and rewatering. Plant Signal. Behav. 5, 649–654.
- [9] Patanè, C.; Saita, A.; Sortino, O. (2013). Comparative effects of salt and water stress on seed germination and early embryo growth in two cultivars of sweet sorghum. J. Agron. Crop Sci. 199, 30–37.
- [10] Prasad, P. V. V.; Djanaguiraman, M.; Jagadish, S. V. K.; Ciampitti, I. A. (2019). Drought and high temperature stress and traits associated with tolerance. In Sorghum: A State of the Art and Future Perspectives; Ciampitti, I. A., Prasad, P. V. V., Eds.; ASA, CSSA, SSSA: Madison, WI, USA, Volume 58, pp. 245–265.
- [11] Earl, H. J.; Davis, R. F. (2003). Effect of drought stress on leaf and whole canopy radiation use efficiency and yield of maize. Agron. J. 95, 688–696.
- [12] Djanaguiraman, M.; Prasad, P. V. V.; Ciampitti, I. A.; Talwar, H. S. (2020). Impact of abiotic stress on sorghum physiology. In: Sorghum in the 21st Century: Food—Fodder—Feed—Fuel for a Rapidly Changing World; Springer Nature: Singapore, pp. 157–188.
- [13] Khan Awais, Valpuri Sovero and Dorcus Gemenet, (2016). Genome-assisted Breeding for Drought Resistance. Current Genomics 17 (4): 330-342.
- [14] Bibi A, Sadaqat HA, Tahir MHN, Akram HM. (2012). Screening of Sorghum (Sorghum bicolor Var Moench) for Drought Tolerance at Seedling Stage in Polyethylene Glycol. Journal of Animal and Plant Sciences, 22 (3): 671-678.
- [15] Abdel-Ghany S, Ullah F, Ben-Hur A, Reddy A. (2020). Transcriptome Analysis of Drought- Resistant and Drought-Sensitive Sorghum (Sorghum bicolor) Genotypes in Response to PEGInduced Drought Stress. International Journal of Molecular Sciences 27 (772).



- [16] Fracasso A, Trindade L, Amaducci S. (2016a). Drought stress tolerance strategies revealed by RNA-Seq in two sorghum genotypes with contrasting WUE. *BMC Plant Biology*, 16: 115.
- [17] Winchell F, Stevens CJ, Murphy C, Champion L, Fuller Dorian Q. (2017). Evidence for Sorghum Domestication in Fourth Millennium BC Eastern Sudan: Spikelet Morphology from Ceramic Impressions of the Butana Group. *Current Anthropology* 58, 673-683.
- [18] Henderson AN, Crim PM, Cumming JR, Hawkins JS. (2020). Phenotypic and physiological responses to salt exposure in Sorghum reveal diversity among domesticated landraces. *American Journal of Botany* 107, 983-992.
- [19] Swigonova Z, Lai J, Ma J, Ramakrishna W, Llaca V, Bennetzen J, Messing J. (2004). Close split of sorghum and maize genome progenitors. *Genome Research*, 1916 1923.
- [20] Qadir M, Bibi A, Tahir MHN, Saleem M, Sadaqat H. (2015). Screening of sorghum (*Sorghum bicolor* L.) genotypes under various levels of drought stress. *Maydica* 60.
- [21] Fracasso A, Trindade L, Amaducci S. (2016b). Drought tolerance strategies highlighted by two Sorghum bicolor races in a dry-down experiment. *Journal of Plant Physiology* 190, 1-14.
- [22] Khalid, W., Ali, A., Arshad, M. S., Afzal, F., Akram, R., Siddeeq, A., et al. (2022). Nutrients and bioactive compounds of Sorghum bicolor L. Used to prepare functional foods: A review on the efficacy against different chronic disorders. *Int. J. Food Prop.* 25 (1), 1045–1062. doi: 10.1080/10942912.2022.2071293.
- [23] Rao, B. D. (2019). "Sorghum value chain for food and fodder security," in breeding sorghum for diverse end uses (Woodhead Publishing), 409–419.
- [24] Dicko MH, Gruppen, H, Traore AS, Van Berkel WJH and Voragen AGJ (2005). Evaluation of the effect of germination on phenolic compounds and antioxidant activities in sorghum varieties. *Journal of Agricultural and Food Chemistry* 53: 2581–2588.
- [25] Rooney LW. (2007). Food and nutritional quality of sorghum and millet. *INTSORMIL 2007 Annual Report*, Lincoln, Nebraska, USA, pp 91–93.
- [26] Shin SI, Choi HJ, Chung KM, Hamaker BR, Park KH and Moon TW. (2004). Slowly digestible starch from debranched waxy sorghum starch: Preparation and properties. *Cereal Chemistry* 81: 404–408.
- [27] Awika JM and Rooney LW. (2004). Sorghum phytochemicals and their potential aspects on human health. *Phytochemistry* 65: 1199–1221.
- [28] Dykes L, Rooney LW, Waniska RD and Rooney WL (2005). Phenolic compounds and antioxidant activity of sorghum grains of varying genotypes. *Journal of Agricultural and Food Chemistry* 53: 6813–6818.
- [29] Cha-um, S., Yooyongwech, S. & Supaibulwatana, K. (2012) Water-deficit tolerant classification in mutant lines of indica rice. *Scientia Agricola*, 69 (2), 135–141.
- [30] Németh, M., Janda, T., Horváth, E., Páldi, E. & Szalai, G. (2002) Exogenous salicylic acid increases polyamine content but may decrease drought tolerance in maize. *Plant Science*, 162 (4), 569–574.
- [31] Ngara, R., Goche, T., Swanevelder, D. Z. H., and Chivasa, S. (2021). Sorghum's whole-plant transcriptome and proteome responses to drought stress: a review. *Life* 11, 704. doi: 10.3390/life11070704
- [32] Fracasso A, Trindade LM, Amaducci S (2016). Drought stress tolerance strategies revealed by RNA-Seq in two sorghum genotypes with contrasting WUE. *BMC Plant Biol.* <https://doi.org/10.1186/s12870-016-0800-x>
- [33] Zhang DF, ZengG TR, Liu XY, Gao CX, Li YX, Li CH, Song YC, shi YS, Wang TY, Yu LI (2019). Transcriptomic profiling of sorghum leaves and roots responsive to drought stress at the seedling stage. *Journal of Integrative Agriculture*, 18 (9), pp. 1980-1995.
- [34] Sukumaran S, Li X, Li X, Zhu C, Bai G, Perumal R, Tuinstra MR, Prasad PV, Mitchell SE, Tesso TT (2016). QTL mapping for grain yield, flowering time, and stay-green traits in sorghum with genotyping-by-sequencing markers. *Crop Sci* 56 (4): 1429–1442. <https://doi.org/10.2135/cropsci2015.02.0097>
- [35] Chaves MM, Flexas J, Pinheiro C (2009). Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Ann Bot* 103 (4): 5 51–560. <https://doi.org/10.1093/aob/mcn125>
- [36] Vadez V, Krishnamurthy L, Hash C, Upadhyaya H, Borrell A (2011b). Yield, transpiration efficiency, and water-use variations and their interrelationships in the sorghum reference collection. *Crop Pasture Sci* 62 (8): 645–655. <https://doi.org/10.1071/CP110 07>
- [37] Lopez JR, Erickson JE, Munoz P, Saballos A, Felderhoff TJ, Vermerris W (2017). QTLs associated with crown root angle, stomatal conductance, and maturity in Sorghum. *Plant Genome*. <https://doi.org/10.3835/plantgenome2016.04.0038>
- [38] Hadebe, S. T.; Modi, A. T.; Mabhaudhi, T. (2017). Drought tolerance and water use of cereal crops: A focus on sorghum as a food security crop in sub-Saharan Africa. *J. Agron. Crop Sci.* 203, 177–191. [CrossRef]
- [39] Ndlovu E, van Staden J, Maphosa M (2021) Morpho-physiological effects of moisture, heat and combined stresses on Sorghum bicolor [Moench (L.)] and its acclimation mechanisms. *Plant Stress*. <https://doi.org/10.1016/j.stress.2021.100018>
- [40] Queiroz MS, Oliveira CE, Steiner F, Zuffo AM, Zoz T, Vendruscolo EP, Silva MV, Mello BFFR, Cabra RC, Menis FT (2019). Drought stresses on seed germination and early growth of maize and sorghum. *Journal of Agricultural Science*, 11 (2), pp. 310-318.
- [41] Ramu VS, Swetha TN, Sheela SH, Babitha CK, Rohini S, Reddy MK, Tuteja N, Reddy CP, Prasad TG, Udayakumar M (2016). Simultaneous expression of regulatory genes associated with specific drought - adaptive traits improves drought adaptation in peanut. *Plant Biotechnology Journal*, 14 (3), pp. 1008-1020.
- [42] Fadoul HE, El Siddig MA, Abdalla AWH, El Hussein AA (2018). Physiological and proteomic analysis of two contrasting Sorghum bicolor genotypes in response to drought stress. *Australian Journal of Crop Science*, 12 (9), pp. 1543-1551.
- [43] Sarshad A, Talei D, Torabi M, Rafiei F, Nejatkhah P (2021). Morphological and biochemical responses of Sorghum bicolor (L.) Moench under drought stress. *SN Applied Sciences*, 3 (1), pp. 1-12.

- [44] Sabadin PK., Malosetti M, Boer MP, Tardin FD, Santo FG, Guimaraes CT, Gomide RL, Andrade CLT, Albuquerque PEP, Caniato FF, Mollinari M (2012). Studying the genetic basis of drought tolerance in sorghum by managed stress trials and adjustments for phenological and plant height differences. *Theoretical and Applied Genetics*, 124 (8), pp. 1389-1402.
- [45] Hattori T, Inanaga S, Araki H, An P, Morita S, Luxová M, Lux A. (2005). Application of silicon enhanced drought tolerance in *Sorghum bicolor*. *Physiologia Plantarum*, 123 (4), pp. 459-466.
- [46] Kapanigowda MH, Perumal R, Djanaguiraman M, Aiken RM, Tesso T, Prasad PV, Little CR, (2013). Genotypic variation in sorghum [*Sorghum bicolor* (L.) Moench] exotic germplasm collections for drought and disease tolerance. *Springer Plus*, 2 (1), pp. 1-13.
- [47] Manjarrez-Sandoval PEDRO, González-Hernández VA, Mendoza-Onofre LE, Engleman E. M (1989). Drought stress effects on the grain yield and panicle development of sorghum. *Canadian Journal of Plant Science*, 69 (3), pp. 631-641.
- [48] Batista - Silva W, Heinemann B, Rugen N, Nunes - Nesi A, Araújo WL, Braun HP, Hildebrandt TM (2019). The role of amino acid metabolism during abiotic stress release. *Plant, cell & environment*, 42 (5), pp. 1630-1644.
- [49] Yu SM, Lo SF, Ho THD (2015). Source-sink communication: regulated by hormone, nutrient, and stress cross-signaling. *Trends Plant Sci* 20 (12): 844-857. <https://doi.org/10.1016/j.tplants.2015.10.009>
- [50] Impa SM, Perumal R, Bean SR, Sunoj VJ, Jagadish SK. (2019). Water deficit and heat stress induced alterations in grain physico-chemical characteristics and micronutrient composition in field grown grain sorghum. *Journal of Cereal Science*, 86, pp. 124-131.
- [51] Stagnari F, Galieni A, Pisante M (2016). Drought stress effects on crop quality. In: *Water Stress and Crop Plants*. pp 375-392. <https://doi.org/10.1002/9781119054450.ch23>
- [52] Ashok Badigannavar, Niaba Teme, Antonio Costa de Oliveira, Guying Li, Michel Vaksman, Vivian Ebeling Viana, T. R. Ganapathi and Fatma Sarsu, (2018). Physiological, genetic and molecular basis of drought resilience in sorghum [*Sorghum bicolor* (L.) Moench]. *Indian Journal for Plant Physiology*. <https://doi.org/10.1007/s40502-018-0416-2>
- [53] Craufurd, P. Q.; Qi, A. (2001). Photothermal adaptation of sorghum (*Sorghum bicolor*) in Nigeria. *Agric. For. Meteorol.* 108, 199-211.
- [54] Barnabas, B.; Jager, K.; Feher, A. (2008). The effect of drought and heat stress on reproductive processes in cereals. *Plant Cell Environ.* 31, 11-38.
- [55] Shavrukov Y, Kurishbayev A, Jatayev S, Shvidchenko V, Zotova L, Koekemoer F, de Groot S, Soole K and Langridge P, (2017). Early Flowering as a Drought Escape Mechanism in Plants: How Can It Aid Wheat Production?. *Front. Plant Sci.* 8: 1950. doi: 10.3389/fpls.2017.01950.
- [56] Hatfield JL and Dold C., (2019). Water-Use Efficiency: Advances and Challenges in a Changing Climate. *Front. Plant Sci.* 10: 103. doi: 10.3389/fpls.2019.00103.
- [57] Tesfamichael Abraha, Stephen Mwangi Githiri, Remmy Kasili, Woldeamlak Araia, Aggrey Bernard Nyende, (2015). Genetic Variation among Sorghum (*Sorghum bicolor* L. Moench) Landraces from Eritrea under Post-Flowering Drought Stress Conditions. *American Journal of Plant Sciences*, 6, 1410-1424. <http://dx.doi.org/10.4236/ajps.2015.69141>
- [58] Velazco JG, Jordan DR, Mace ES, Hunt CH, Malosetti M and van Eeuwijk FA, (2019). Genomic Prediction of Grain Yield and Drought-Adaptation Capacity in Sorghum Is Enhanced by Multi-Trait Analysis. *Front. Plant Sci* 10: 997. doi: 10.3389/fpls.2019.00997.
- [59] Kassahun, B.; Bidinger, F. R.; Hash, C. T.; Kuruvashetti, M. S. (2009). Stay-green expression in early generation sorghum (*Sorghum bicolor* (L.) Moench) QTL introgression lines. *Euphytica*, 172, 351-362.
- [60] Gano, B.; Dembele, J. S. B.; Tovignan, T. K.; Sine, B.; Vadez, V.; Diouf, D.; Audebert, A., (2021). Adaptation Responses to Early Drought Stress of West Africa Sorghum Varieties. *Agronomy* 11 (443): 1-21. <https://doi.org/10.3390/agronomy11030443>
- [61] Mahalakshmi, V.; Bidinger, F. R. (2002). Evaluation of putative stay-green sorghum germplasm lines. *Crop Sci.* 42, 965-974.
- [62] Passioura, J. B. (2012). Phenotyping for drought tolerance in grain crops: When is it useful to breeders? *Funct. Plant Biol.* 39, 851-859.
- [63] Turner, N. C.; Begg, J. E.; Tonnet, M. L. (1978). Osmotic adjustment of sorghum and sunflower crops in response to water deficits and its influence on the water potential at which stomata close. *Funct. Plant Biol.* 5, 597-608.
- [64] Blum A. (2005). Drought resistance, water-use efficiency, and yield potential—are they compatible, dissonant, or mutually exclusive? *Australian Journal of Agricultural Research* 56, 1159.
- [65] Ogawa A, Yamauchi A. (2006). Root Osmotic Adjustment under Osmotic Stress in Maize Seedlings 1. Transient Change of Growth and Water Relations in Roots in Response to Osmotic Stress. *Plant Production Science* 9, 27-38.
- [66] Assefa, Y.; Staggenborg, S. A.; Prasad, P. V. V. (2010). Grain sorghum water requirement and responses to drought stress: A review. *Crop. Manag.* 2010, doi: 10.1094/CM-1109-01-RV.
- [67] Turner NC, Jones MM. (1980). Turgor Maintenance by Osmotic Adjustment: A Review and Evaluation. In: Turner, N. C. and Kramer, P. J., Eds., *Adaptation of Plants to Water and High Temperature Stress*, Wiley & Sons, New York, 87-103.
- [68] Blum A. (2017). Osmotic adjustment is a prime drought stress adaptive engine in support of plant production: Osmotic adjustment and plant production. *Plant, Cell & Environment* 40, 4-10.
- [69] Chaves M, Costa J, Zarrouk O, Pinheiro O, Lopes C, Pereira J. (2016). Controlling stomatal aperture in semi-arid regions – The dilemma of saving water or being cool. *Plant Science* 251, 54-64.
- [70] Marcinska I, Czyczyło-Mysza I, Skrzypek E, Filek M, Grzesiak S, Grzesiak M, Janowiak F, Hura T, Dziurka M, Dziurka K, Nowakowska A, Quarrie S. (2012). Impact of osmotic stress on physiological and biochemical characteristics in drought-susceptible and drought resistant wheat genotypes. *Acta Physiologiae Plantarum* 35, 451-461.

- [71] Rodrigues, J., Inze, D., Nelissen, H., and Saibo, N. J. M. (2019). Source-sink regulation in crops under water deficit. *Trends Plant Sci.* 24, 652–663. doi: 10.1016/j.tplants.2019.04.005.
- [72] Xue D, Zhang X, Lu X, Chen G and Chen Z-H, (2017). Molecular and Evolutionary Mechanisms of Cuticular Wax for Plant Drought Tolerance. *Front. Plant Sci.* 8: 621. doi: 10.3389/fpls.2017.00621.
- [73] Elango, D., Xue, W. & Chopra, S., (2020). Genome wide association mapping of epi-cuticular wax genes in Sorghum bicolor. *Physiol Mol Biol Plants* 26, 1727–1737. <https://doi.org/10.1007/s12298-020-00848-5>
- [74] Sajeevan R. S., Parvathi M. S. and Nataraja K. N., (2017). Leaf wax trait in crops for drought and biotic stress tolerance: regulators of epicuticular wax synthesis and role of small RNAs. *Ind J plant physio.* 22,434-447 <https://doi.org/10.1007/s40502-017-0333-9>
- [75] Amtmann, A., Bennett, M. J., and Henry, A. (2022). Root phenotypes for the future. *Plant Cell Environ.* 45 (3), 595–601. doi: 10.1111/pce.14269.
- [76] Ruben Rufo, Silvio Salvi, Conxita Royo and Jose Miguel Soriano, 2020. Exploring the Genetic Architecture of Root-Related Traits in Mediterranean Bread Wheat Landraces by Genome-Wide Association Analysis. *Agronomy*10: 613doi: 10.3390/agronomy10050613.
- [77] Fromm, H. (2019). Root plasticity in the pursuit of water. *Plants-Basel.* 8 (7), 236. doi: 10.3390/plants8070236.
- [78] Gupta, A., Rico-Medina, A., and Cano-Delgado, A. I. (2020). The physiology of plant responses to drought. *Science* 368 (6488), 266–269. doi: 10.1126/science.aaz7614.
- [79] Siddiqui, M. N., Leon, J., Naz, A. A., and Ballvora, A. (2021). Genetics and genomics of root system variation in adaptation to drought stress in cereal crops. *J. Exp. Bot.* 72 (4), 1007–1019. doi: 10.1093/jxb/eraa487.
- [80] Kang, J., Peng, Y., and Xu, W. (2022). Crop root responses to drought stress: molecular mechanisms, nutrient regulations, and interactions with microorganisms in the rhizosphere. *Int. J. Mol. Sci.* 23, 9310. doi: 10.3390/ijms23169310.
- [81] Fenta, B. A.; Beebe, S. E.; Kunert, K. J.; Burrridge, J. D.; Barlow, K. M.; Lynch, P. J. (2014). Fiel phenotyping of soybean roots for drought stress tolerance. *Agronom* 4, 418–435.
- [82] Comas Louise H., Becker Steven R., Cruz Von Mark V., Byrne Patrick F. and Dierig David A., (2013). Root traits contributing to plant productivity under drought. *Frontiers in Plant Science* 4: 442 doi: 10.3389/fpls.2013.00442.
- [83] Allah Wasaya, Xiyang Zhang, Qin Fang and Zongzheng Yan, (2018). Root Phenotyping for Drought Tolerance: A Review. *Agronomy* 8: 241; doi: 10.3390/agronomy8110241.
- [84] Vega, R. C., Villagra, P. E., and Greco, S. A. (2020). Different root strategies of perennial native grasses under two contrasting water availability conditions: implications for their spatial distribution in desert dunes. *Plant Ecol.* 221 (7) 633–646. doi: 10.1007/s11258-020-01038-9.
- [85] Lynch, J. P. (2022). Harnessing root architecture to address global challenges. *Plant J.* 109 (2), 415–431. doi: 10.1111/tbj.15560.
- [86] Chen, X.; Wu, Q.; Gao, Y.; Zhang, J.; Wang, Y.; Zhang, R.; Zhou, Y.; Xiao, M.; Xu, W.; Huang, R., (2020). The role of deep roots in sorghum yield production under drought conditions. *Agronomy* 10, 611.
- [87] Hasan, M. M.; Rafii, M. Y.; Ismail, M. R.; Mahmood, M.; Rahim, H. A.; Alam, M. A.; Ashkani, S.; Malek, M. A.; Latif, M. A. (2015). Marker assisted backcrossing: A useful method for rice improvement. *Biotechnol. Biotechnol. Equip.* 29, 237–254. [CrossRef] [PubMed]
- [88] Sanchez, A. C.; Subudhi, P. K.; Rosenow, D. T.; Nguyen, H. T. (2002). Mapping QTLs associated with drought resistance in sorghum [*Sorghum bicolor* (L.) Moench]. *Plant Mol. Biol.* 48, 713–726. [CrossRef] [PubMed]
- [89] Crasta, O. R.; Xu, W. W.; Rosenow, D. T.; Mullet, J.; Nguyen, H. T. (1999). Mapping of post-flowering drought resistance traits in grain sorghum: Association between QTLs influencing premature senescence and maturity. *Mol. Genet. Genom.* 262, 579–588. [CrossRef] [PubMed]
- [90] Harris, K. R. (2007) Genetic Analysis of the Sorghum bicolor Stay-Green Drought Tolerance Trait. Ph. D. Thesis, Texas A&M University, College Station, TX, USA.
- [91] Hash, C. T.; Raj, A. G. B.; Lindup, S.; Sharma, A.; Beniwal, C. R.; Folkertsma, R. T.; Mahalakshmi, V.; Zerbini, E.; Blummel, M. (2003). Opportunities for marker- assisted selection (MAS) to improve the feed quality of crop residues in pearl millet and sorghum. *Field Crop. Res.* 84, 79–88.
- [92] Baloch, F. S.; Alsaleh, A.; Shahid, M. Q.; Çiftçi, V.; de Miera, L. E. S.; Aasim, M.; Nadeem, M. A.; Akta,s, H.; Özkan, H.; Hatipoğlu, R.(2017). A Whole Genome DArTseq and SNP Analysis for Genetic Diversity Assessment in Durum Wheat from Central Fertile Crescent. *PLoS ONE*, 12, e0167821. [CrossRef]
- [93] Mwamahonje, A.; Eleblu, J. S. Y.; Ofori, K.; Feyissa, T.; Deshpande, S.; Garcia-Oliveira, A. L.; Bohar, R.; Kigoni, M.; Tongoona, P. (2021). Introgression of QTLs for Drought Tolerance into Farmers’ Preferred Sorghum Varieties. *Agriculture*, 11, 883. [CrossRef]
- [94] Subudhi, P. K.; Rosenow, D. T.; Nguyen, H. T. (2007). Quantitative trait loci for the stay green trait in sorghum (*Sorghum bicolor* L. Moench): Consistency across genetic backgrounds and environments. *Theor. Appl. Genet.* 101, 733–741. [CrossRef]
- [95] Menz, M. A.; Klein, R. R.; Mullet, J. E.; Obert, J. A.; Unruh, N. C.; Klein, P. E. (2002). A high-density genetic map of *Sorghum bicolor* (L.) Moench based on 2926 AFLP, RFLP and, SSR markers. *Plant Mol. Biol.* 48, 483–499. [CrossRef]
- [96] Reddy, R. N.; Madhusudhana, R.; Mohan, S. M.; Chakravarthi, D. V.; Mehtre, S. P.; Seetharama, N.; Patil, J. V. (2013). [*Sorghum bicolor* (L.) Moench] Mapping QTL for grain yield and other agronomic traits in post-rainy sorghum. *Theor. Appl. Genet.* 126, 1921–1939. [CrossRef]
- [97] Kholova, J.; Deshpande, S.; Madhusudhana, R.; Blummel, M.; Borrell, A.; Hammer, G. (2019). Improving Post-Rainy Sorghum Varieties to Meet the Growing Grain and Fodder Demand in India-Phase 2; Australian Centre for International Agricultural Research: Canberra, Australia, p. 29.
- [98] Kadam, S. R.; Fakrudin, B. (2017). Marker assisted pyramiding of root volume QTLs to improve drought tolerance in rabi sorghum. *Res. Crop.* 18, 683–692. [CrossRef]

- [99] Takeda Sh, and Matsuoka M, (2008). Genetic approaches to crop improvement: Responding to environmental and population changes. *Nature Reviews Genetics*. 9 (6): 44-57. doi: 10.1038/nrg2342. [PubMed]
- [100] Kamal, N. M.; Gorafi, Y. S. A.; Ghanim, A. M. A. (2017). Performance of Sorghum stay-green introgression lines under post-flowering drought. *Int. J. Plant Res.* 7, 65–74.
- [101] Sakiyama, N. S.; Ramos, H. C. C.; Caixeta, E. T.; Pereira, M. G. (2014). Plant breeding with marker-assisted selection in Brazil. *Crop Breed. Appl. Biotechnol.* 14, 54–60. [CrossRef]
- [102] Kebede, H.; Subudhi, P. K.; Rosenow, D. T.; Nguyen, H. T. (2001). Quantitative trait loci influencing drought tolerance in grain sorghum [*Sorghum bicolor* (L.) Moench]. *Theor. Appl. Genet.* 103, 266–276. [CrossRef]
- [103] Gorthy, S.; Narasu, L.; Gaddameedi, A.; Sharma, H. C.; Kotla, A.; Deshpande, S. P.; Are, A. K. (2017). Introgression Introgression of Shoot Fly [*Atherigona soccata* (L.) Moench] Resistance QTLs into Elite Post-rainy Season Sorghum Varieties Using Marker Assisted Backcrossing (MABC). *Front. Plant Sci.* 8, 1494. [CrossRef]
- [104] Ongom, P. O. (2016). Association Mapping of Gene Regions for Drought Tolerance and Agronomic Traits in Sorghum. Ph. D. Thesis, Purdue University, West Lafayette, IN, USA.
- [105] Rida Fatima Ahmed, Muhammad Irfan, Hafiz Abdullah Shakir, Muhammad Khan and Lijing Chen, (2020). Engineering drought tolerance in plants by modification of transcription and signalling factors. *Biotechnology & Biotechnological Equipment*, 34 (1): 781-789, <https://doi.org/10.1080/13102818.2020.1805359>
- [106] Sharmila Polumahanthi, Sarada M ani N., Sudhakar Pola, Dora S. V. V. S. N. and Nageswara Rao S. (2014). Tissue Culture, Molecular and Genetic Approaches to Sorghum Crop Improvement – A Review. *Indian Journal of Plant Sciences* Vol. 4 Pp. 97-113.
- [107] Elkonin L. A., O. N. Nosova and J. V. Italianskaya, (2012). Agrobacterium-Mediated Genetic Transformation of Sorghum Using Tissue Culture-Based and Pollen Mediated Approaches. *Journal of Stress Physiology & Biochemistry*, Vol. 8 No. 3.
- [108] Bray, E. A. (1993). Molecular responses to water deficit. *Plant Physiol.* 103, 103 1040. doi: 10.1104/pp.103.4.1035.
- [109] Bray, E. A. (1997). Plant responses to water deficit. *Trends Plant Sci.* 2, 48–54. doi: 10.1016/S1360-1385(97)82562-9.
- [110] Xiong, L., and Zhu, J. (2002). Molecular and genetic aspects of plant responses to osmotic stress. *Plant Cell Environ.* 25, 131–139. doi: 10.1046/j.1365-3040.2002.00782.x.
- [111] Farooq, M., Wahid, A., Kobayashi, N., Fujita, D., and Basra, S. M. A. (2009). Plant drought stress: effects, mechanisms and management. *Agron. Sustain. Dev.* 29, 185– 212. doi: 10.1051/agro:2008021.
- [112] Basu, S., Ramegowda, V., Kumar, A., and Pereira, A. (2016). Plant adaptation to drought stress. *F1000 Res.* 5, 1554. doi: 10.12688/f1000research.7678.1.
- [113] Zhang, J., Jiang, F., Shen, Y., Zhan, Q., Bai, B., Chen, W., et al. (2019a). Transcriptome analysis reveals candidate genes related to phosphorus starvation tolerance in sorghum. *BMC Plant Biol.* 19, 306. doi: 10.1186/s12870-019-1914-8.
- [114] Wang, T., Ren, Z., Liu, Z., Feng, X., Guo, R., Li, B., et al. (2014). SbHKT1; 4, a member of the high-affinity potassium transporter gene family from sorghum bicolor, functions to maintain optimal Na<sup>+</sup> /K<sup>+</sup> balance under Na<sup>+</sup> stress. *J. Integr. Plant Biol.* 56, 315–332. doi: 10.1111/jipb.12144.
- [115] Li, H., Li, Y., Ke, Q., Kwak, S. S., Zhang, S., and Deng, X. (2020). Physiological and differential proteomic analyses of imitation drought stress response in *Sorghum bicolor* root at the seedling stage. *Int. J. Mol. Sci.* 21, 9174. doi: 10.3390/ijms21239174.