

Research Article

Evaluation of Growth Promotion and Pathogenicity of Endophytic Fungi from the Root of *Chenopodium Quinoa* Willd

Victor Paco Pérez^{1,*} , Crespo Melgar Carla Fabiana² ,
Fernando Pacasa Quisbert¹ 

¹Technical University of Oruro, Faculty of Agricultural and Natural Sciences, Oruro, Plurinational State of Bolivia

²Mayor University of San Andrés, Faculty of Pharmaceutical and Biological Sciences, La Paz, Plurinational State of Bolivia

Abstract

Symbiotic associations between endophytic fungi and *Chenopodium quinoa* have been reported to be beneficial for increasing tolerance to drought and soil salinity, being this the only crop grown in the Southern Altiplano of Bolivia. The symbiotic interaction of endophytic fungi was investigated, considering their detrimental and mutualistic effects. 38 strains of endophytic fungi from the fungal collection of the Faculty of Agronomy were used. The percentage of germination and root length were measured in vitro (4 days). The percentage of emergence (7 days) and plant height (14 days) in pots were also evaluated. The fungi that did not impair germination were strains VP42, VP44, *Alternaria* sp. VP37, *Fusarium* sp. VP05 and *Fusarium* sp. VP30. Fungi that stimulated a significant increase in radicle longitudinal growth were *Fusarium* sp. VP35, *Alternaria* sp. VP37 and strain VP18. Fungi that caused a high mortality rate during the emergence phase were strain VP01, *Alternaria* sp. VP15, *Fusarium* sp. VP02, *Fusarium* sp. VP07, *Fusarium* sp. VP08, *Fusarium* sp. VP12, *Fusarium* sp. VP23 and *Fusarium* sp. VP36. The endophytic strains *Alternaria* sp. VP37, *Fusarium* sp. VP35 and strain VP18 stimulated superior seedling growth. The present research work reveals that some endophytic fungi of the *Alternaria* and *Fusarium* genera can behave as pathogens during the germination stage, while others have the function of promoting quinoa growth.

Keywords

Alternaria, *Fusarium*, Endophytic Fungus, Pathogenesis, Quinoa

1. Introduction

Quinoa (*Chenopodium quinoa* Willd.), is an annual dicotyledonous plant belonging to Amaranthaceae family [1]. Currently, it is cultivated in more than 125 countries around the world, due to its genetic diversity of seeds preserved by generations of farmers in the Andes [2]. It is the only crop

adapted for thousands of years to the adverse conditions of the Southern Altiplano of Bolivia [3, 4].

Plant-fungus interactions date back 400-500 million years [5-7]. Plants do not thrive isolated, as constantly interact closely with various microorganisms, mainly bacteria and

*Corresponding author: laramvico@gmail.com (Victor Paco Pérez)

Received: 24 November 2024; **Accepted:** 23 December 2024; **Published:** 9 January 2025



Copyright: © The Author(s), 2025. Published by Science Publishing Group. This is an **Open Access** article, distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

fungi [8-11]. Their symbiotic relationship with plants attracts attention for its benefits in agriculture.

Fungi benefits to their host, existing a wide variety of plant-fungal symbiotic interactions [12-15]. Favorable interactions, for example, allow plants to adapt to various ecosystems [16-19]. For quinoa, endophytic fungi (EF) perform various functions that improve its resistance to adverse conditions, mitigating abiotic stress [20, 21].

Associations can be beneficial [8, 22-25], for example, promoting the adaptation and growth of plants in extreme conditions [26]. Most plants interact with microorganisms [27, 28], EF have been reported in plants that inhabit the Arctic, Antarctic, desert, ocean, tropical forests and agricultural fields [29-33]. Growth promoting fungi can protect against plant pathogens by increasing tolerance to abiotic stress, such as drought, salinity and high temperatures through the production of secondary metabolites and hydrolytic enzymes [34-36]. This study evaluated the influence of EF inoculated in roots and its effects on germination, emergence, root development and growth promotion in quinoa. Understanding the effect of applying EF in agriculture can offer innovative solutions to increase quinoa crop productivity by mitigating the conditions of the Southern Altiplano of Bolivia.

2. Materials and Methods

2.1. Study Area

The trials were carried out at the Plant Science Laboratory of the Faculty of Agronomy of the Technical University of Oruro. The HE strains used in this study were partially identified and are part of the mycological collection of the Faculty of Agronomy, whose specific details are presented in Table 1.

2.2. Fungal Cultivation and Biomass Production

For *in vitro* evaluation, axenic culture of EF were prepared in Potato Dextrose Agar (PDA) medium at 25 °C for 14 days. Subsequently, mycelia discs (5 mm) of EF were collected [37] to evaluate their promoting or pathogenic effect on quinoa germination. For pot tests, EF biomass was produced in 250 mL flasks inoculating 10 mycelia discs in 50 g of sugarcane bagasse.

2.3. In Vitro Germination and Root Length

Seeds of quinoa (Pandela rosada variety) were disinfested according to González-Teuber et al. [21]. *In vitro* evaluation of EF effect on quinoa germination and root length, consisted of applying a mycelia disk in a 100 mm Petri dish with Murashige and Skoog (MS) medium (38 HE with 3 replicates + uninoculated Petri dishes) along with 10 disinfested quinoa seeds located around the disk. Petri dishes were incubated at

16 °C for 4 days, with daily monitoring. Finally, germination (%) and root length (cm) were evaluated after the incubation period [38, 39].

Table 1. Endophytic fungi in the current study.

Specimens	Strain code	Host	Place
Alternaria sp.	VP15	Quinoa root	Southern Altiplano - Bolivia
Alternaria sp.	VP24	Quinoa root	Southern Altiplano - Bolivia
Alternaria sp.	VP37	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP12	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP11	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP16	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP02	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP21	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP22	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP23	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP03	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP30	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP35	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP36	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP39	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP40	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP41	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP43	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP05	Quinoa root	Southern Altiplano - Bolivia
Fusarium sp.	VP06	Quinoa root	Southern Altiplano - Bolivia

Specimens	Strain code	Host	Place
<i>Fusarium</i> sp.	VP07	Quinoa root	Southern Altiplano - Bolivia
<i>Fusarium</i> sp.	VP08	Quinoa root	Southern Altiplano - Bolivia
<i>Penicillium</i> sp.	VP32	Quinoa root	Southern Altiplano - Bolivia
<i>Penicillium</i> sp.	VP38	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP01	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP14	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP18	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP19	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP20	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP28	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP29	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP31	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP33	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP34	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP04	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP42	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP44	Quinoa root	Southern Altiplano - Bolivia
Unknown	VP09	Quinoa root	Southern Altiplano - Bolivia

2.4. Pots Assay: Emergence and Plant Length

Inoculation of EF in pot experiments consisted of: i) 15 g of sugarcane bagasse colonized with the fungus was placed at the bottom, ii) 600 g of sterile soil (collected from fallow quinoa plots in Southern Altiplano, commonly sandy) into 1 L plastic pot (38 pots with 3 replicates + uninoculated pots). Then, all pots were covered with sterile plastic bags until sowing. 10 disinfested quinoa seeds were sown at the surface and covered again with sterile plastic bags until germination. The pots were kept at room temperature (14 °C) with relative humidity of 37%. The emergence percentage (%) was evaluated at 7 days and the plant length (cm) at 14 days after sowing.

2.5. Statistical Analysis

A complete randomized design (CRD) was used for the *in vitro* evaluation and a complete randomized block design (RBD) for the pots assay. Analysis of variance (ANOVA) was determined using R 4.2.3 software (R Development Core 2023) and Tukey test was used post-hoc.

3. Results

In vitro assay. There were significant differences in germination of quinoa seeds subjected to 38 EF ($p < 0.05$). EF such as *Alternaria* sp. VP37, *Alternaria* sp. VP24, *Fusarium* sp. VP05, *Fusarium* sp. VP30, strains VP44 and VP42 applied to quinoa seeds promoted 100% of germination. In contrast, EF as *Alternaria* sp. VP15, *Fusarium* sp. VP12, *Fusarium* sp. VP23, *Fusarium* sp. VP02, *Fusarium* sp. VP08, *Fusarium* sp. VP07, strain VP01 and *Fusarium* sp. VP36 inhibited the germination completely, but *Fusarium* sp. VP21, strain VP14, *Fusarium* sp. VP43, *Fusarium* sp. VP39, strain VP31, *Fusarium* sp. VP22 and *Fusarium* sp. VP16 inhibited 93.33% of germination. Other strains showed intermediate values (Figure 1). On the other hand, there were significant differences in root length among the 38 EF ($p < 0.05$). Growth promoting fungi based on longitudinal development of roots were *Fusarium* sp. VP35, *Alternaria* sp. VP37 and strain VP18 that reached 49.85, 49.73 and 48.85 mm respectively, followed by *Alternaria* sp. VP24, *Fusarium* sp. VP05, strains VP29, VP42 and VP44 that promote relatively similar quinoa roots varying from 46.43 to 46.12 mm. In contrast, the strain VP14 and *Fusarium* sp. VP43 affected root development with 23.28 and 21.98 mm, respectively (Figures 2, 3).




Figure 1. Quinoa germination can be influenced by inoculation with endophytic fungi.




Figure 2. Root development of quinoa can be influenced by inoculation with endophytic fungi.




Figure 3. In vitro test: Endophytic fungi effect on quinoa root length. (A) Fungus observed to be beneficial as it promotes root growth, (B) Fungus has a moderately detrimental effect on root development, and (C) certain endophytic fungi are detrimental to root development.

Emergence and height of potted plants. Fully emerged cotyledons were statistically different in the presence of EF ($p < 0.05$). The endophytic fungi that did not cause damage were strains VP18, VP19, VP29, VP42, VP44, *Fusarium* sp. VP35, *Alternaria* sp. VP37, *Fusarium* sp. VP05, *Alternaria* sp. VP24 and *Fusarium* sp. VP30, which only limited emergence by 3.33%. EF that presented less emergence of cotyledons and caused high mortality were *Alternaria* sp. VP15, *Fusarium* sp. VP07, *Fusarium* sp. VP08, *Fusarium* sp. VP02, *Fusarium* sp. VP12, *Fusarium* sp. VP36, strain VP01 and *Fusarium* sp. VP23 with 96.67% (Figure 4).

Plant height was statistically significant ($p < 0.05$). Endophytic strains *Alternaria* sp. VP37, *Fusarium* sp. VP35 and strain VP18 promoted higher plant development reaching 55.98, 54.95 and 53.58 mm respectively, followed by strains VP44, VP42, VP19 and *Fusarium* sp. VP05 with developments 44.91, 44.86, 41.52 and 41.62 mm, respectively. *Fusarium* sp. VP03, *Fusarium* sp. VP06, as well as strains

VP04, VP31 and VP33, showed a development similar to that of plants free of infection by endophytic fungi (control). Other endophytic strains had negative effects on plant height compared to the control (Figures 5, 6).

4. Discussion

Some EF from quinoa roots have positive effects on germination, root length, emergence and plant height. However, the vast majority caused detrimental effects including death. Several studies have described the positive effect of the association between EF and several crops [32, 40–46]. The HES that were isolated from the quinoa root of the Atacama Desert, had their positive effect on the morphological and physiological characteristics of quinoa by inoculation of *Talaromyces minioluteus* and *Penicillium murcianum* under drought and salinity conditions [20, 46].




Figure 4. Effect of inoculation with endophytic fungi in the emergence phase of quinoa.




Figure 5. The effect of inoculation with endophytic fungi on the height growth of quinoa plants.

EF from quinoa root such as *Alternaria* sp. VP37, *Fusarium* sp. VP05, *Fusarium* sp. VP30, strains VP44 and VP42 promoted 100% of *in vitro* (MS and PDA medium) germination compared to other endophytes. Moreover, under controlled conditions in pots assay, strains *Fusarium* sp. VP35, *Fusarium* sp. VP05, *Alternaria* sp. VP37, strains VP18, VP19,

VP42 and VP44 promoted 100% of plant emergence (Figure 4). The symbiosis between EF and plants can increase or reduce certain phytohormones, for example, by secreting indole acetic acid (IAA) and gibberellic acid (GA) that affect longitudinal growth of roots and phyllosphere [47].




Figure 6. Impact of endophytic fungi inoculation on quinoa growth. In (A), a positive boost in development is evident due to the mutualistic strain. On the other hand, in (B), it is observed that the endophytic fungus has a moderate detrimental effect, while in (C), it is shown that this strain is highly detrimental due to its pathogenic behavior.

Three EF that promoted greatest root elongation were *Fusarium* sp. VP35, *Alternaria* sp. VP37 and strain VP18,

reaching a plant height of 55.98, 54.95 and 53.58 mm (Figure 5). These results reveal that fungi of the *Fusarium* genus may

promote plant growth, probably by producing IAA, nitrogen fixation, phosphate solubilization and production of siderophores such as *F. annulatum* and *F. proliferatum* [48]. Nevertheless, the relationship between plant-fungus can be beneficial and/or pathogenic [49], for example, those symbionts that do not cause any disease symptoms can colonize host plants tissues and there are benefits for both parties [50]. In contrast, harmful aspects are described to interfere in transport of nutrients and water [51], abnormal metabolism in different stages of plant growth [52]. These negative effects can be controlled by other EF [53], for example, *A. fumigatus* that has the ability to control pathogens by producing aflatoxins [54]. Based upon the above described, the results of the initial vegetative development in this study showed that EF can be harmful and/or beneficial on the development of quinoa (Figure 6). For example, the strains *Alternaria* sp. VP37, *Fusarium* sp. VP35 and strain VP18 promoted seedling growth, but *Fusarium* sp. VP02 and *Fusarium* sp. VP07 are the strains that affected longitudinal elongation of the seedlings. In root development, 25.92% of EF promoted growth and the rest restricted elongation. These results are similar to those of other authors such as [55] who reported that there are pathogenic fungi of the genus *Penicillium*. In contrast, there are studies that highlighted how EF stimulate the root growth to promote nutrients absorption [44], in exchange the plants synthesize several hydrolytic enzymes such as xylanase, pectinase, cellulase and proteinase that favor the fungi to access plant tissues [32]. Understanding the dynamics of interaction between endophytic fungi and crop plants is essential to take advantage of the positive effects and mitigate the risks of potential pathogenic effects. That is essential for the development of agricultural management strategies that optimize the use of endophytic fungi in the improvement of crops such as quinoa, especially under abiotic stress conditions, such as drought and salinity.

5. Conclusion

Some strains of EF significantly promote key processes such as germination, root growth, emergence and height development of quinoa plants. However, other strains exert inhibitory effects on these processes, which could be detrimental.

EF with growth-promoting capacity represent a resource with great potential for implementation as biostimulants in sustainable agricultural systems. However, it is imperative to carry out a rigorous and precise selection of strains before proceeding with their application in field conditions.

Abbreviations

EF	Endophytic Fungi
MS	Murashige and Skoog
PDA	Potato Dextrose Agar

IAA	Indole Acetic Acid
GA	Gibberellic Acid

Source of Financing

This research did not receive external funding.

Acknowledgments

We would like to thank Edwin Marcelo Gonzales Torrico for managing the microbiology laboratory of the Agricultural Engineering Degree of the Faculty of Agricultural and Natural Sciences of the Technical University of Oruro. We would also like to thank the National Institute of Agricultural and Forestry Innovation. We would also like to thank the Peasant Academic Unit of Tiahuanacu of the Bolivian Catholic University "San Pablo." We would also like to thank quinoa producers from different municipalities of the Southern Altiplano of Bolivia.

Ethical Considerations

This work was carried out according to the recommendations of related authors.

Research Limitations

There were no limitations in the present research.

Author Contributions

Victor Paco Pérez, conceived the idea and wrote the initial draft of the manuscript, Carla Crespo Melgar and Fernando Pacasa Quisbert, edited and revised the manuscript.

Conflicts of Interest

The authors declare no conflicts of interest.

References

- [1] A. Vega-Gálvez, M. Miranda, J. Vergara, E. Uribe, L. Puente, y E. A. Martínez, «Nutrition facts and functional potential of quinoa (*Chenopodium quinoa* Willd.), an ancient Andean grain: a review», *J. Sci. Food Agric.*, vol. 90, n.º 15, pp. 2541-2547, dic. 2010, <https://doi.org/10.1002/jsfa.4158>
- [2] D. Bazile, «Global Trends in the Worldwide Expansion of Quinoa Cultivation», *Biol. Life Sci. Forum*, vol. 15, n.º 1, p. 13, 2023, [En línea]. Disponible en: <https://doi.org/10.3390/blsf2023025013>
- [3] D. Bazile, S.-E. Jacobsen, y A. Verniau, «The Global Expansion of Quinoa: Trends and Limits», *Front. Plant Sci.*, vol. 7, p. 622, 2016, <https://doi.org/10.3389/fpls.2016.00622>

- [4] M. Walsh-Dilley, «Resilience compromised: Producing vulnerability to climate and market among quinoa producers in Southwestern Bolivia», *Glob. Environ. Chang.*, vol. 65, p. 102165, 2020, <https://doi.org/10.1016/j.gloenvcha.2020.102165>
- [5] F. Lutzoni *et al.*, «Contemporaneous radiations of fungi and plants linked to symbiosis», *Nat. Commun.*, vol. 9, n.º 1, p. 5451, 2018, <https://doi.org/10.1038/s41467-018-07849-9>
- [6] C. Strullu-Derrien, M. A. Selosse, P. Kenrick, y F. M. Martin, «The origin and evolution of mycorrhizal symbioses: from palaeomycology to phylogenomics», *New Phytol.*, vol. 220, n.º 4, pp. 1012-1030, 2018, <https://doi.org/10.1111/nph.15076>
- [7] N. Vigneron, G. V. Radhakrishnan, y P. M. Delaux, «What have we learnt from studying the evolution of the arbuscular mycorrhizal symbiosis?», *Curr. Opin. Plant Biol.*, vol. 44, pp. 49-56, 2018, <https://doi.org/10.1016/j.pbi.2018.02.004>
- [8] M. A. Hassani, P. Durán, y S. Hacquard, «Microbial interactions within the plant holobiont», *Microbiome*, pp. 6-58, 2018, https://doi.org/10.1007/978-1-4419-9863-7_100630
- [9] J. Mercado-Blanco, «Life of Microbes Inside the Plant», en *Principles of Plant-Microbe Interactions: Microbes for Sustainable Agriculture*, B. Lugtenberg, Ed. Springer, 2015, pp. 25-32. <https://doi.org/10.1007/978-3-319-08575-3>
- [10] W. K. Mousa y M. N. Raizada, «The diversity of anti-microbial secondary metabolites produced by fungal endophytes: an interdisciplinary perspective», *Front. Microbiol.*, vol. 4, p. 65, 2013, <https://doi.org/10.3389/fmicb.2013.00065>
- [11] Z.-L. Yuan, C.-L. Zhang, y F.-C. Lin, «Role of Diverse Non-Systemic Fungal Endophytes in Plant Performance and Response to Stress: Progress and Approaches», *J. Plant Growth Regul.*, vol. 29, n.º 1, pp. 116-126, 2010, <https://doi.org/10.1007/s00344-009-9112-9>
- [12] P. Bonfante y A. Genre, «Mechanisms underlying beneficial plant - Fungus interactions in mycorrhizal symbiosis», *Nat. Commun.*, p. 1: 48, 2010, <https://doi.org/10.1038/ncomms1046>
- [13] F. A. Decunta, L. I. Pérez, D. P. Malinowski, M. A. Molina-Montenegro, y P. E. Gundel, «A Systematic Review on the Effects of Epichloë Fungal Endophytes on Drought Tolerance in Cool-Season Grasses», *Front. Plant Sci.*, vol. 12, p. 644731, 2021, <https://doi.org/10.3389/fpls.2021.644731>
- [14] K. M. Thambugala, D. A. Daranagama, A. J. L. Phillips, S. D. Kannangara, y I. Promputtha, «Fungi vs. Fungi in Biocontrol: An Overview of Fungal Antagonists Applied Against Fungal Plant Pathogens», *Front. Cell. Infect. Microbiol.*, vol. 10, n.º November, p. 604923, 2020, <https://doi.org/10.3389/fcimb.2020.604923>
- [15] P. Trivedi, J. E. Leach, S. G. Tringe, T. Sa, y B. K. Singh, «Plant-microbiome interactions: from community assembly to plant health», *Nat. Rev. Microbiol.*, vol. 18, n.º 11, pp. 607-621, 2020, <https://doi.org/10.1038/s41579-020-0412-1>
- [16] A. E. Fadji y O. O. Babalola, «Exploring the potentialities of beneficial endophytes for improved plant growth», *Saudi J. Biol. Sci.*, vol. 27, n.º 12, pp. 3622-3633, 2020, <https://doi.org/10.1016/j.sjbs.2020.08.002>
- [17] P. R. Hardoim *et al.*, «The Hidden World within Plants: Ecological and Evolutionary Considerations for Defining Functioning of Microbial Endophytes», *Microbiol. Mol. Biol. Rev.*, vol. 79, n.º 3, pp. 293-320, 2015, <https://doi.org/10.1128/mmbr.00050-14>
- [18] K. Selim, «Biology of Endophytic Fungi», *Curr. Res. Environ. Appl. Mycol.*, vol. 2, n.º 1, pp. 31-82, 2012, <https://doi.org/10.5943/cream/2/1/3>
- [19] V. V. Kumar, «Plant Growth-Promoting Microorganisms: Interaction with Plants and Soil», en *Plant, Soil and Microbes*, vol. 1, K. R. Hakeem, M. S. Akhtar, y S. N. A. Abdullah, Eds. Springer International Publishing Switzerland, 2016, pp. 1-16. https://doi.org/10.1007/978-3-319-27455-3_1
- [20] M. González-Teuber, R. A. Contreras, G. E. Zúñiga, D. Barrera, y L. Bascuñán-Godoy, «Synergistic Association With Root Endophytic Fungi Improves Morpho-Physiological and Biochemical Responses of Chenopodium quinoa to Salt Stress», *Front. Ecol. Evol.*, vol. 9, p. 787318, 2022, <https://doi.org/10.3389/fevo.2021.787318>
- [21] M. González-Teuber, C. Vilo, y L. Bascuñán-Godoy, «Molecular characterization of endophytic fungi associated with the roots of Chenopodium quinoa inhabiting the Atacama Desert, Chile», *Genomics Data*, vol. 11, pp. 109-112, 2017, <https://doi.org/10.1016/j.gdata.2016.12.015>
- [22] G. Berg, P. Kusstatscher, A. Abdelfattah, T. Cernava, y K. Smalla, «Microbiome Modulation-Toward a Better Understanding of Plant Microbiome Response to Microbial Inoculants», *Front. Microbiol.*, vol. 12, p. 650610, 2021, <https://doi.org/10.3389/fmicb.2021.650610>
- [23] Y. A. Millet *et al.*, «Innate immune responses activated in Arabidopsis roots by microbe-associated molecular patterns», *Plant Cell*, vol. 22, n.º 3, pp. 973-990, 2010, <https://doi.org/10.1105/tpc.109.069658>
- [24] A. C. Perez, J. S. Rojas, y H. M. Vale, «Biología y perspectiva de microorganismos endófitos asociados a plantas», *Rev. Colomb. Cienc. Anim. - RECIA*, vol. 1, n.º 2, pp. 286-301, 2009, <https://doi.org/10.24188/recia.v1.n2.2009.372>
- [25] A. Zuccaro *et al.*, «Endophytic life strategies decoded by genome and transcriptome analyses of the mutualistic root symbiont *Piriformospora indica*», *PLoS Pathog.*, vol. 7, n.º 10, p. e1002290, 2011, <https://doi.org/10.1371/journal.ppat.1002290>
- [26] I. Khalmuratova, D. H. Choi, H. J. Yoon, T. M. Yoon, y J. G. Kim, «Diversity and plant growth promotion of fungal endophytes in five halophytes from the buan salt marsh», *J. Microbiol. Biotechnol.*, vol. 31, n.º 3, pp. 408-418, 2021, <https://doi.org/10.4014/JMB.2012.12041>
- [27] A. E. Arnold y F. Lutzoni, «Diversity and host range of foliar fungal endophytes: are tropical leaves biodiversity hotspots?», *Ecology*, vol. 88, n.º 3, pp. 541-549, 2007, <https://doi.org/10.1890/05-1459>

- [28] S. Kusari, S. P. Pandey, y M. Spiteller, «Untapped mutualistic paradigms linking host plant and endophytic fungal production of similar bioactive secondary metabolites», *Phytochemistry*, vol. 91, pp. 81-87, 2012, <https://doi.org/10.1016/j.phytochem.2012.07.021>
- [29] N. M. Chauhan, A. D. Gutama, y A. Aysa, «Endophytic fungal diversity isolated from different agro-ecosystem of Enset (Ensete ventricosum) in Gedeo zone, SNNPRS, Ethiopia», *BMC Microbiol.*, vol. 19, n.º 1, p. 172, 2019, [En línea]. Disponible en: <https://doi.org/10.1186/s12866-019-1547-y>
- [30] J. G. Harrison y E. A. Griffin, «The diversity and distribution of endophytes across biomes, plant phylogeny and host tissues: how far have we come and where do we go from here?», *Environ. Microbiol.*, vol. 22, n.º 6, pp. 2107-2123, 2020, <https://doi.org/10.1111/1462-2920.14968>
- [31] L. Jin *et al.*, «Diversity and Biocontrol Potential of Culturable Endophytic Fungi in Cotton», *Front. Microbiol.*, vol. 12, p. 698930, 2021, <https://doi.org/10.3389/fmicb.2021.698930>
- [32] K. L. Rana *et al.*, «Endophytic microbes: biodiversity, plant growth-promoting mechanisms and potential applications for agricultural sustainability», *Antonie Van Leeuwenhoek*, vol. 113, n.º 8, pp. 1075-1107, 2020, <https://doi.org/10.1007/s10482-020-01429-y>
- [33] R. E. Sánchez-Fernández *et al.*, «Hongos endófitos: fuente potencial de metabolitos secundarios bioactivos con utilidad en agricultura y medicina», *Tip Rev. Espec. en Ciencias Químico-Biológicas*, vol. 16, n.º 2, pp. 132-146, 2013, [https://doi.org/10.1016/s1405-888x\(13\)72084-9](https://doi.org/10.1016/s1405-888x(13)72084-9)
- [34] J. A. Castillo, G. Conde, M. Claros, y N. Ortuño, «Diversity of cultivable microorganisms associated with Quinoa (Chenopodium quinoa) and their potential for plant growth-promotion», *Bionatura*, vol. 7, n.º 2, p. 61, 2022, <https://doi.org/10.21931/RB/2022.07.02.61>
- [35] J. O. de Carvalho, V. Broll, A. H. S. Martinelli, y F. C. Lopes, «Endophytic fungi: positive association with plants», en *Molecular Aspects of Plant Beneficial Microbes in Agriculture*, V. Sharma, R. Salwan, y L. K. Tawfeeq Al-Ani, Eds. India: Academic Press, 2020, pp. 321-332. [En línea]. Disponible en: <https://doi.org/10.1016/B978-0-12-818469-1.00026-2>
- [36] N. Ortuño, J. A. Castillo, y M. Claros, «Symbiotic Native Microorganisms of Quinoa in the Bolivian Altiplano», en *Biology and Biotechnology of Quinoa*, A. Varma, Ed. Springer Nature Singapore, 2021, pp. 131-151. [En línea]. Disponible en: https://doi.org/10.1007/978-981-16-3832-9_7
- [37] P. W. Crous *et al.*, «Fusarium: more than a node or a foot-shaped basal cell», *Stud. Mycol.*, vol. 98, p. 100116, 2021, <https://doi.org/10.1016/j.simyco.2021.100116>
- [38] M. R. Panuccio, S. E. Jacobsen, S. S. Akhtar, y A. Muscolo, «Effect of saline water on seed germination and early seedling growth of the halophyte quinoa», *AoB Plants*, vol. 6, p. plu047, 2014, <https://doi.org/10.1093/aobpla/plu047>
- [39] A. Naureen, F. ul H. Nasim, M. S. Choudhary, M. Ashraf, F. M. W. Grundler, y A. S. S. Schleker, «A new endophytic fungus CJAN1179 isolated from the Cholistan desert promotes lateral root growth in Arabidopsis and produces IAA through tryptophan-dependent pathway», *Arch. Microbiol.*, vol. 204, n.º 3, p. 181, 2022, <https://doi.org/10.1007/s00203-022-02768-2>
- [40] N. C. Baron, A. de Souza Pollo, y E. C. Rigobelo, «Purpureocillium lilacinum and Metarhizium marquandii as plant growth-promoting fungi», *PeerJ*, n.º 8, p. e9005, 2020, <https://doi.org/10.7717/peerj.9005>
- [41] G. Berg, «Plant-microbe interactions promoting plant growth and health: Perspectives for controlled use of microorganisms in agriculture», *Appl. Microbiol. Biotechnol.*, vol. 84, n.º 1, pp. 11-18, 2009, <https://doi.org/10.1007/s00253-009-2092-7>
- [42] L. R. Jaber y J. Enkerli, «Fungal entomopathogens as endophytes: can they promote plant growth?», *Biocontrol Sci. Technol.*, vol. 27, n.º 1, pp. 28-41, 2017, <https://doi.org/10.1080/09583157.2016.1243227>
- [43] A. L. Khan, J. Hussain, A. Al-Harrasi, A. Al-Rawahi, y I.-J. Lee, «Endophytic fungi: resource for gibberellins and crop abiotic stress resistance», *Crit. Rev. Biotechnol.*, vol. 35, pp. 62-74, 2015.
- [44] N. C. Baron y E. C. Rigobelo, «Endophytic fungi: a tool for plant growth promotion and sustainable agriculture», *Mycology*, vol. 13, n.º 1, pp. 39-55, 2022, <https://doi.org/10.1080/21501203.2021.1945699>
- [45] L. Yan, J. Zhu, X. Zhao, J. Shi, C. Jiang, y D. Shao, «Beneficial effects of endophytic fungi colonization on plants», *Appl. Microbiol. Biotechnol.*, vol. 103, n.º 8, pp. 3327-3340, 2019, <https://doi.org/10.1007/s00253-019-09713-2>
- [46] M. González-Teuber, A. Urzuá P. Plaza, y L. Bascuñán-Godoy, «Effects of root endophytic fungi on response of Chenopodium quinoa to drought stress», *Plant Ecol.*, vol. 219, pp. 231-240, 2018, <https://doi.org/10.1007/s11258-017-0791-1>
- [47] L. Bilal *et al.*, «Plant growth promoting endophytic fungi Asprgillus fumigatus TS1 and Fusarium proliferatum BRL1 produce gibberellins and regulates plant endogenous hormones», *Symbiosis*, vol. 76, pp. 117-127, 2018, [En línea]. Disponible en: <https://doi.org/10.1007/s13199-018-0545-4>
- [48] S. Roy, A. Sarma, S. Paul, D. K. Jha, y K. Tayung, «Plant growth-promoting traits and activation of defense enzymes in traditional rice variety by fungal endophytes isolated from seeds of indigenous rice cultivars of Northeast India», *South African J. Bot.*, vol. 160, pp. 483-492, 2023, [En línea]. Disponible en: <https://doi.org/10.1016/j.sajb.2023.07.041>
- [49] N. A. Elhamouly *et al.*, «The hidden power of secondary metabolites in plant-fungi interactions and sustainable phytoremediation», *Front. Plant Sci.*, vol. 13, p. 1044896, 2022, <https://doi.org/10.3389/fpls.2022.1044896>
- [50] P. Tiwari y H. Bae, «Endophytic Fungi: Key Insights, Emerging Prospects, and Challenges in Natural Product Drug Discovery», *Microorganisms*, vol. 10, p. 360, 2022, <https://doi.org/10.3390/microorganisms10020360>

- [51] S. E. Smith, E. Facelli, S. Pope, y F. A. Smith, «Plant performance in stressful environments: interpreting new and established knowledge of the roles of arbuscular mycorrhizas», *Plant Soil*, vol. 326, pp. 3-20, 2010, <https://doi.org/10.1007/s11104-009-9981-5>
- [52] H. A. Bruns, «Controlling Aflatoxin and Fumonisin in Maize by Crop Management», *J. Toxicol.*, vol. 22, n.º 2-3, pp. 153-173, 2003, <https://doi.org/10.1081/TXR-120024090>
- [53] M. Kottb, T. Gigolashvili, D. K. Grofkinsky, y B. Piechulla, «Trichoderma volatiles effecting Arabidopsis: from inhibition to protection against phytopathogenic fungi», *Front. Microbiol.*, vol. 6, p. 995, 2015, <https://doi.org/10.3389/fmicb.2015.00995>
- [54] A. M. Abdelaziz, D. A. El-Wakil, M. S. Attia, O. M. Ali, H. AbdElgawad, y A. H. Hashem, «Inhibition of *Aspergillus flavus* Growth and Aflatoxin Production in *Zea mays* L. Using Endophytic *Aspergillus fumigatus*», *J. Fungi*, vol. 8, n.º 5, p. 482, 2022, <https://doi.org/10.3390/jof8050482>
- [55] J. P. Louw y L. Korsten, «Pathogenic *Penicillium* spp. on apple and pear», *Plant Dis.*, vol. 98, n.º 5, pp. 590-598, 2014, <https://doi.org/10.1094/PDIS-07-13-0710-RE>