

# Some Upper Bounds of Maximum $E$ -Eigenvalues of Uniform Hypergraphs

Hongyu Zhang<sup>\*</sup>, Feng Fu, Caoji Yin

School of Mathematics and Statistics, Qinghai Normal University, Xining, China

**Email address:**

1103403129@qq.com (Hongyu Zhang)

<sup>\*</sup>Corresponding author

**To cite this article:**

Hongyu Zhang, Feng Fu, Caoji Yin. Some Upper Bounds of Maximum  $E$ -Eigenvalues of Uniform Hypergraphs. *Mathematics and Computer Science*. Vol. 8, No. 2, 2023, pp. 51-56. doi: 10.11648/j.mcs.20230802.13

**Received:** February 25, 2023; **Accepted:** March 16, 2023; **Published:** March 28, 2023

**Abstract:** A hypergraphs, as a generalization of a general graph, is often used as an effective tool to describe complex structures in discrete mathematics, computer science and other fields. Hypergraph theory and related parameters of hypergraph are important research topics in hypergraph theory. In particular, the problem of spectral extremum of graphs has been widely concerned. This problem originates from the problem proposed by Brualdi and Solheid in 1986. That is to find the upper and lower bounds of spectral radius of a given graph class and characterize the polar graph that reaches the upper and lower bounds. Let  $H$  be a uniform hypergraph. Let  $A(H)$  be the adjacency tensor of  $H$ . In this work, by using Perron-Frobenius theorem, Hölder’s inequality and inequality of arithmetic and geometric means, we establish some upper bounds for the maximum  $E$ -eigenvalue of a uniform hypergraph instead of the degrees of vertices and edge number of hypergraph  $H$ . In addition, we characterize the extremal hypergraphs that reach the upper bounds.

**Keywords:** Uniform Hypergraphs, Adjacency Tensor, Maximal  $E$ -Eigenvalue, Degree

## 1. Introduction

Denote the set  $\{1, 2, \dots, n\}$  by  $[n]$ . Let  $H = (V(H), E(H))$  be a hypergraph with  $n$  vertexes and  $m$  edges, where  $V(H) = \{v_1, v_2, \dots, v_n\}$  and  $E(H) = \{e_1, e_2, \dots, e_m\}$  represent the vertex set and edge set of the hypergraph  $H$ , respectively. If every edge  $e_i = \{v_{i_1}, v_{i_2}, \dots, v_{i_l}\}$  in a hypergraph  $H$  satisfies  $|e_i| = l = r$ , then a hypergraph  $H$  is called  $r$ -uniform hypergraph, where  $i_j \in [n], j = 1, 2, \dots, l$  and  $i = 1, 2, \dots, m$ . Let  $d_i$  be the degree of the vertex  $v_i$ . If  $d_i = d$  is satisfied for any  $i \in [n]$ , then the hypergraph  $H$  is called a  $d$ -regular hypergraph. Let  $\Delta$  and  $\delta$  be the minimum degree and the maximum degree of hypergraph  $H$ , respectively. Tensor's definition is as follows.

Definition 1 [10]. Given any positive integers  $r, n$ , a tensor  $\mathcal{T}$  of order  $r$  and dimension  $n$  is defined by a

multidimensional array of element  $t_{i_1, i_2, \dots, i_r}$ , where  $i_1, i_2, \dots, i_r \in [n]$ ,  $t_{i_1, i_2, \dots, i_r} \in C$ . Obviously, a tensor  $\mathcal{T}$  of order  $r$  and dimension  $n$  has  $n^r$  elements.

The definition of tensor product is given, which is a generalization of matrix product [8].

Definition 2 [10, 4]. Let  $\mathcal{A} \in C^{n_1 \times n_2 \times \dots \times n_m}$ ,  $\mathcal{B} \in C^{n_2 \times n_3 \times \dots \times n_{r+1}}$  be two tensors of orders  $m$  and  $r$ , respectively. Then the product  $\mathcal{C} = \mathcal{A}\mathcal{B}$  of tensors  $\mathcal{A}$  and  $\mathcal{B}$  is a tensor of order  $(m-1)(r-1)+1$  with the  $(i, \alpha_1, \alpha_2, \dots, \alpha_{m-1})$ -th entry is

$$C_{i, \alpha_1, \alpha_2, \dots, \alpha_{m-1}} = \sum_{i_2, i_3, \dots, i_m \in [n_2]} \mathcal{A}_{i, i_2, i_3, \dots, i_m} \mathcal{B}_{i_2, \alpha_1, \dots, \alpha_{m-1}, i_m}$$

where  $i \in [n_1], \alpha_1, \alpha_2, \dots, \alpha_{m-1} \in [n_3] \times [n_4] \times \dots \times [n_{r+1}]$ .

Definition 3 [10, 14]. Let  $\mathcal{A} \in C^{n_1 \times n_2 \times \dots \times n_m}$ ,

$$\mathcal{B} \in C^{n_2 \times n_3 \times \dots \times n_{r+1}}$$

be two tensors of orders  $m$  and  $r$ , respectively.  $\mathcal{T}\mathbf{x}^{r-1}$  is a  $n$ -dimension complex vector with the  $i$ -th entry:

$$\left(\mathcal{T}\mathbf{x}^{r-1}\right)_i = \sum_{i_2, i_3, \dots, i_r=1}^r t_{ii_2 \dots i_r} x_{i_2} x_{i_3} \dots x_{i_r}, i \in [n].$$

$\mathcal{T}\mathbf{x}^r$  is a complex number:

$$\mathcal{T}\mathbf{x}^r = \sum_{i_1, i_2, \dots, i_r=1}^r t_{i_1 i_2 \dots i_r} x_{i_1} x_{i_2} \dots x_{i_r}, i \in [n].$$

In 2005, Qi and Lim proposed the definition of tensor eigenvalues, respectively [2, 3]. If there exists  $\lambda \in C$  and a non-zero vector  $\mathbf{x} \in C^n$  such that

$$\mathcal{T}\mathbf{x}^{r-1} = \lambda \mathbf{x}^{[r-1]},$$

then  $\lambda$  is the eigenvalue of tensor  $\mathcal{T}$ , and  $\mathbf{x}$  is the corresponding eigenvector related with  $\lambda$  of  $\mathcal{T}$ , where

$$\mathbf{x}^{[r-1]} = \left(x_1^{r-1}, x_2^{r-1}, \dots, x_n^{r-1}\right)^T.$$

In 2012, Cooper and Dutle defined the adjacency tensor  $\mathcal{A}(H)$  of a  $r$ -uniform hypergraph  $H$  [1].

Definition 4 [1]. Let  $H = (V(H), E(H))$  be a uniform hypergraph with  $n$  vertexes. The adjacency tensor  $\mathcal{A}(H)$  of hypergraph  $H$  is a tensor of order  $r$  and dimension  $n$ , whose  $(i_1, i_2, \dots, i_r)$ -th entry is

$$a_{i_1 i_2 \dots i_r} = \begin{cases} \frac{1}{(r-1)!}, & \text{if } \{i_1, i_2, \dots, i_r\} \in E(H) \\ 0, & \text{otherwise} \end{cases}$$

Qi et al. gave the definition of  $E$ -eigenvalue of a uniform hypergraph [11].

Definition 5 [11]. Given a tensor  $\mathcal{A}$  of order  $r$  and dimension  $n$ . Let  $\lambda \in C$  and  $\mathbf{x} \in C^n$ , if

$$\lambda \mathbf{x} = \mathcal{A}\mathbf{x}^{r-1}, \mathbf{x}\mathbf{x}^T = 1,$$

then  $\mathbf{x}$  is the  $E$ -eigenvector of tensor  $\mathcal{A}$  related with its eigenvalue  $\lambda$ .

Lemma 1 [6, 7, 13]. (Perron-Frobenius theorem) Let  $\mathcal{A}$  be a non-negative weakly irreducible tensor of order  $r$  and dimension  $n$ , where  $r, n \geq 2$ . Let  $\lambda$  is the maximum eigenvalue of  $\mathcal{A}$ . Then  $\lambda$  is the only eigenvalue that satisfies the corresponding eigenvector is positive.

In 2009, Bulo and Pelillo published the first paper on the spectrum of a hypergraph, and used the concept of  $H$ -eigenvalue of a tensor introduced by Qi in 2005 [12]. By

the upper bound of clique numbers of graphs derived from Sós and Straus, Motzkin-Straus theorem is extended to specific hypergraphs [15]. In particular, Bulo and Pelillo extended and improved the spectral boundary introduced by Wilf, and established a link between the cluster number and the spectral hypergraph theory [12]. In 2012, Cooper and Dutle systematically studied hypergraph theory through adjacency tensor, and extended the basic results of graph theory to hypergraphs [1]. Kang and Liu et al. gave some upper bounds on the compactness of the eigenvalues of hypergraphs as follows [5].

$$\rho(H) \leq \max_{1 \leq i \leq n} \left( \frac{1}{r-1} \sum_{\{v_{i_1}, v_{i_2}, \dots, v_{i_r}\} \in E(H)} \left( d_{i_2}^{r-1} + d_{i_3}^{r-1} + \dots + d_{i_r}^{r-1} \right) \right)^{\frac{r-1}{r}}.$$

Equality holds if and only if the hypergraph  $H$  is a regular hypergraph.

Besides, Yi gave an upper bound of  $H$ -eigenvalue of a hypergraph [9].

$$\rho(H) \leq \max_{\{v_{i_1}, v_{i_2}, \dots, v_{i_r}\} \in E(H)} \sqrt[2(r-1)]{d_{i_1}^{r-1} d_{i_2}^{r-1} d_{i_3}^{r-1} \dots d_{i_r}^{r-1}}.$$

In this work, several upper bounds of the maximal  $E$ -eigenvalues of uniform hypergraphs are established in terms of their degrees by using Perron-Frobenius theorem and Holder inequality. In addition, the extremal hypergraphs are characterized.

## 2. Main Results

Let  $E(H(v_1))$  be all edges containing vertex  $v_1$ . Write

$$\sum_{\{v_1, v_{i_2}, \dots, v_{i_r}\} \in E(H(v_1))} x_{i_2} x_{i_3} \dots x_{i_r} = \sum_{e \in E(H(v_1))} x^e$$

for short.

Theorem 1. Suppose  $H$  is a  $r$ -uniform hypergraph with  $n$  vertices. Then

$$\lambda \leq \Delta r^{\frac{r(r-2)}{4(r-1)}}.$$

Equality holds if and only if the hypergraph  $H$  has only one hyperedge, which contains all vertices.

*Proof* According to Lemma 1,

$$\mathbf{x} = (x_1, x_2, \dots, x_n)^T, x_i > 0, i \in [n].$$

Let the entry  $x_1$  corresponding to vertex  $v_1$  be the largest entry in the eigenvector  $\mathbf{x}$ . Suppose edge  $e_1 = \{v_1, v_{i_2}, \dots, v_{i_r}\}$  is the edge with the largest product of eigenvector entries. According to the characteristic equation, we have

$$\lambda x_1 = \sum_{e \in E(H(v_1))} x^e \leq d_1 x_2 x_3 \dots x_r, \tag{1}$$

$$\begin{cases} \lambda x_2 = \sum_{e \in E(H(v_2))} x^e \leq d_2 x_1^{r-1}, \\ \lambda x_3 = \sum_{e \in E(H(v_3))} x^e \leq d_3 x_1^{r-1}, \\ \vdots \\ \lambda x_r = \sum_{e \in E(H(v_r))} x^e \leq d_r x_1^{r-1}. \end{cases} \tag{2}$$

Multiply the corresponding  $r-1$  inequalities (1), then

$$\lambda^{r-1} (x_2 x_3 \dots x_r) \leq (d_2 d_3 \dots d_r) x_1^{(r-1)^2}. \tag{3}$$

If the both sides of inequality (2) are  $r-1$  power at the same time, then

$$\lambda^{r-1} x_1^{r-1} \leq d_1^{r-1} (x_2 x_3 \dots x_r)^{r-1}. \tag{4}$$

Multiply the two sides of inequality (3) and (4), respectively, then

$$\lambda^{2(r-1)} x_1^{r-1} (x_2 x_3 \dots x_r) \leq d_1^{r-1} (d_2 d_3 \dots d_r) x_1^{(r-1)^2} (x_2 x_3 \dots x_r)^{r-1}. \tag{5}$$

By  $d_i \leq \Delta, i \in [n]$  and (5), we obtain

$$\lambda^{2(r-1)} \leq \Delta^{2(r-1)} x_1^{(r-2)^2} \left( x_2^2 x_3^2 \dots x_r^2 \right)^{\frac{r-2}{2}}. \tag{6}$$

From the mean inequality and  $\mathbf{x}\mathbf{x}^T = 1$ , it arrives

$$\lambda^{2(r-1)} \leq \Delta^{2(r-1)} \left( \frac{1}{r} \right)^{\frac{r(r-2)}{2}} x_1^{(r-2)^2}. \tag{7}$$

Since  $\mathbf{x}\mathbf{x}^T = 1, x_1^2 \leq 1$ , we get

$$\lambda^{2(r-1)} \leq \Delta^{2(r-1)} \left( \frac{1}{r} \right)^{\frac{r(r-2)}{2}}. \tag{8}$$

Therefore,

$$\lambda \leq \Delta r^{\frac{r(r-2)}{4(r-1)}}. \tag{9}$$

□

Let  $f(r) = r^{\frac{r(r-2)}{4(r-1)}}$  be a function on  $r$ . When  $r \geq 3$ ,

$\frac{df}{dr} < 0$ . That is,  $f(r)$  is a decreasing function with  $r$ .

When  $r$  reaches the maximum value, the corresponding hypergraph is only one hyperedge with  $n$  vertexes (see Figure 1). Meanwhile,  $x_1 = x_2 = \dots = x_n$ . If  $r = 2$ , then the

hypergraph  $H$  is a general graph satisfying  $\lambda \leq \Delta$ .




Figure 1. Hypergraph  $H_0$ .

Theorem 2. Let  $H$  be a  $r$ -uniform hypergraph with  $n$  vertices. Then

$$\text{Equation Section (Next)} \lambda \leq m r^{\frac{1-r}{2}}.$$

Proof By Lemma 1, we have

$$\mathbf{x} = (x_1, x_2, \dots, x_n)^T, x_i > 0, i \in [n].$$

According to the characteristic equation, then

$$\lambda x_i = \sum_{\{v_i, v_{i_2}, v_{i_3}, \dots, v_{i_r}\} \in e_i \subseteq E(H)} x_{i_2} x_{i_3} \dots x_{i_r}. \tag{10}$$

From Holder inequality, we get

$$\begin{aligned} \lambda x_i &= \sum_{\{v_i, v_{i_2}, v_{i_3}, \dots, v_{i_r}\} \in e_i \subseteq E(H)} 1 \cdot x_{i_2} x_{i_3} \dots x_{i_r} \\ &\leq d_i^{\frac{1}{r}} \left( \sum_{\{v_i, v_{i_2}, v_{i_3}, \dots, v_{i_r}\} \in e_i \subseteq E(H)} (x_{i_2} x_{i_3} \dots x_{i_r})^{\frac{r}{r-1}} \right)^{\frac{r-1}{r}} \\ &\leq d_i^{\frac{1}{r}} x_i^{-1} \left( \sum_{\{v_i, v_{i_2}, v_{i_3}, \dots, v_{i_r}\} \in e_i \subseteq E(H)} (x_{i_2}^2 x_{i_3}^2 \dots x_{i_r}^2)^{\frac{r}{2(r-1)}} \right)^{\frac{r-1}{r}}. \end{aligned} \tag{11}$$

Then,

$$\lambda x_i^2 \leq d_i^{\frac{1}{r}} \left( \sum_{\{v_i, v_{i_2}, v_{i_3}, \dots, v_{i_r}\} \in e_i \subseteq E(H)} (x_{i_2}^2 x_{i_3}^2 \dots x_{i_r}^2)^{\frac{r}{2(r-1)}} \right)^{\frac{r-1}{r}}. \tag{12}$$

As  $\mathbf{x}\mathbf{x}^T = 1, x_1^2 + x_2^2 + \dots + x_n^2 = 1$ . For any  $r$  elements  $x_{i_1}, x_{i_2}, \dots, x_{i_r}$ , we have

$$x_{i_1}^2 + x_{i_2}^2 + \dots + x_{i_r}^2 \leq x_1^2 + x_2^2 + \dots + x_n^2 = 1. \tag{13}$$

According to the mean inequality, we obtain

$$\lambda x_i^2 \leq d_i^{\frac{1}{r}} \left( \sum_{\{v_i, v_{i_2}, v_{i_3}, \dots, v_{i_r}\} \in e_i \subseteq E(H)} \left( \frac{1}{r} \right)^{\frac{r^2}{2(r-1)}} \right)^{\frac{r-1}{r}}. \tag{14}$$

Then

$$\lambda x_i^2 \leq d_i \left( \frac{1}{r} \right)^{\frac{r}{2}}. \tag{15}$$

Summing over  $i$  from 1 to  $n$  for (15), it arrives

$$\lambda \leq rm \times r^{-\frac{r}{2}} = mr^{1-\frac{r}{2}}. \quad (16) \quad \lambda \leq \max_{1 \leq i \leq n} \left( \frac{1}{r-1} \sum_{\{v_1, v_2, \dots, v_r\} \in E(H)} \left( d_{i_2}^{\frac{1}{r-1}} + d_{i_3}^{\frac{1}{r-1}} + \dots + d_{i_r}^{\frac{1}{r-1}} \right) \right)^{\frac{1}{r-1}}.$$

□

Theorem 3. Let  $H$  be a  $r$ -uniform hypergraph with  $n$  vertices. Then

Equality holds if and only if the hypergraph  $H$  is regular. Equation Section (Next).

*Proof* According to the characteristic equation, we have

$$\lambda x_i = \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in e_i \subseteq E(H)} x_{i_2} x_{i_3} \dots x_{i_r}. \quad (17)$$

By Holder inequality, we get

$$\begin{aligned} \lambda x_i &= \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(H)} 1 \cdot x_{i_2} x_{i_3} \dots x_{i_r} \\ &\leq d_i^{\frac{1}{r}} \left( \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(H)} (x_{i_2} x_{i_3} \dots x_{i_r})^{\frac{r}{r-1}} \right)^{\frac{r-1}{r}}. \end{aligned} \quad (18)$$

According to the mean inequality, then

$$\lambda x_i \leq \left( \frac{1}{r-1} \right)^{\frac{r-1}{r}} d_i^{\frac{1}{r}} \left( \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(H)} (x_{i_2}^r + x_{i_3}^r + \dots + x_{i_r}^r) \right)^{\frac{r-1}{r}}. \quad (19)$$

If both sides are  $\frac{r}{r-1}$  power, then

$$(\lambda x_i)^{\frac{r}{r-1}} \leq \frac{1}{r-1} d_i^{\frac{1}{r-1}} \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(H)} (x_{i_2}^r + x_{i_3}^r + \dots + x_{i_r}^r). \quad (20)$$

Then

$$\lambda^{\frac{r}{r-1}} \leq \frac{1}{r-1} d_i^{\frac{1}{r-1}} x_i^{\frac{r-1}{r}} \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(H)} (x_{i_2}^r + x_{i_3}^r + \dots + x_{i_r}^r). \quad (21)$$

Since  $\mathbf{x}\mathbf{x}^T = 1$ ,  $x_1^2 \leq 1$  and  $x_i^{\frac{r-1}{r}} \leq x_i^{-2}$ ,

$$\lambda^{\frac{r}{r-1}} x_i^2 \leq \frac{1}{r-1} d_i^{\frac{1}{r-1}} \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(H)} (x_{i_2}^r + x_{i_3}^r + \dots + x_{i_r}^r). \quad (22)$$

Due to  $\mathbf{x}\mathbf{x}^T = 1$ , we have  $x_1^2 + x_2^2 + \dots + x_n^2 = 1$ .

Summing over  $i$  from 1 to  $n$  for (22), we obtain

$$\begin{aligned} \lambda^{\frac{r}{r-1}} &\leq \frac{1}{r-1} \sum_{i=1}^n \left( \frac{1}{d_i^{\frac{1}{r-1}}} \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(G)} (x_{i_2}^r + x_{i_3}^r + \dots + x_{i_r}^r) \right) \\ &= \frac{1}{r-1} \sum_{i=1}^n \left( \frac{1}{x_i^{\frac{r-1}{r}}} \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(G)} \left( d_{i_2}^{\frac{1}{r-1}} + d_{i_3}^{\frac{1}{r-1}} + \dots + d_{i_r}^{\frac{1}{r-1}} \right) \right) \\ &\leq \frac{1}{r-1} \max_{1 \leq i \leq n} \left( \sum_{\{v_1, v_2, v_3, \dots, v_r\} \in E(G)} \left( d_{i_2}^{\frac{1}{r-1}} + d_{i_3}^{\frac{1}{r-1}} + \dots + d_{i_r}^{\frac{1}{r-1}} \right) \right). \end{aligned} \quad (23)$$

Then

$$\lambda \leq \max_{1 \leq i \leq n} \left( \frac{1}{r-1} \sum_{\{v_{i_1}, v_{i_2}, \dots, v_{i_r}\} \in E(H)} \left( d_{i_2}^{r-1} + d_{i_3}^{r-1} + \dots + d_{i_r}^{r-1} \right) \right)^{\frac{1}{r-1}}. \tag{24}$$

For the above results, equality holds if and only if

$$x_1 = x_2 = \dots = x_n.$$

Obviously, equality holds if and only if the hypergraph  $H$  is regular.  $\square$

A regular hypergraph  $H_1$  is shown in Figure 2.




Figure 2. Hypergraph  $H_1$ .

From theorem 3 and using the maximum and minimum degree of a hypergraph, the following corollary can be obtained.  
Corollary 3.1 Let  $H$  be a  $r$ -uniform hypergraph with  $n$  vertices. Then

$$\lambda \leq \left( \frac{\left( (rm)^{\frac{1}{r-1}} n^{\frac{r-2}{r-1}} + \left( (r-1) \delta^{r-1} \Delta - \Delta^{r-1} \right) - (n-1) \delta^{r-1} \right)}{r-1} \right)^{\frac{1}{r-1}}.$$

*Proof* According to the Power mean inequality, we have

$$\sum_{i=1}^n d_i^{r-1} \leq n^{\frac{r-2}{r-1}} \left( \sum_{i=1}^n d_i \right)^{\frac{1}{r-1}} (rm)^{\frac{1}{r-1}} n^{\frac{r-2}{r-1}}. \tag{25}$$

According to Theorem 3, then

$$(r-1) \lambda^{\frac{r}{r-1}} \leq \max_{1 \leq i \leq n} \left( \sum_{\{v_{i_1}, v_{i_2}, \dots, v_{i_r}\} \in E(H)} \left( d_{i_2}^{r-1} + d_{i_3}^{r-1} + \dots + d_{i_r}^{r-1} \right) \right). \tag{26}$$

Substitute (25) into (26), then

$$(r-1) \lambda^{\frac{r}{r-1}} \leq \max_{1 \leq i \leq n} \left( (rm)^{\frac{1}{r-1}} n^{\frac{r-2}{r-1}} - d_i^{r-1} - (n - (r-1)d_i - 1) \delta^{r-1} \right). \tag{27}$$

So,

$$(r-1)\lambda^{\frac{r}{r-1}} \leq \max_{1 \leq i \leq n} \left( (rm)^{\frac{1}{r-1}} n^{\frac{r-2}{r-1}} + \left( (r-1)\delta^{\frac{1}{r-1}} d_i - d_i^{\frac{1}{r-1}} \right) - (n-1)\delta^{\frac{1}{r-1}} \right). \tag{28}$$

Let  $f(x) = (r-1)\delta^{\frac{1}{r-1}}x - x^{\frac{1}{r-1}}$  be a function of  $x$ . When  $1 \leq \delta \leq x \leq \Delta$ ,  $\frac{df}{dx} < 0$ . That is,  $f(x)$  a decreasing function on  $x$  as  $1 \leq \delta \leq x \leq \Delta$ . When  $x = \Delta$ ,  $f(x) = (r-1)\delta^{\frac{1}{r-1}}\Delta - \Delta^{\frac{1}{r-1}}$ . Therefore,

$$\lambda \leq \left( \frac{(rm)^{\frac{1}{r-1}} n^{\frac{r-2}{r-1}} + \left( (r-1)\delta^{\frac{1}{r-1}}\Delta - \Delta^{\frac{1}{r-1}} \right) - (n-1)\delta^{\frac{1}{r-1}}}{r-1} \right). \tag{29}$$

□

### 3. Conclusion

In this work, using Perron-Frobenius theorem and Holder inequality, some upper bounds of the maximum  $E$ -eigenvalue of hypergraphs are obtained. In fact, spectral radius can effectively characterize the structure of a hypergraph. For example, spectral radius can be used to study the connectivity, diameter and matching number of hypergraphs. Therefore, in the future research, we consider more properties of hypergraph radius, and establish the relationship between the invariant of a hypergraph such as its girth, diameter, matching number, domination number and hypergraph radius.

### Funding

Qinghai Office of Science and Technology (Grant No. 2022-ZJ-T02); NSFC (No. 12261073, No. 11961055); The 111 Project (D20035).

### References

[1] J. Cooper, A. Dutle, Spectra of uniform hypergraphs, *Linear Algebra Appl.* 436 (2012) 3268-3299.

[2] L. H. Lim, Singular values and eigenvalues of tensors, a variational approach, in: *Proceedings of the IEEE International Workshop on Computational Advances in Multi-Sensor Adaptive Processing*, vol. 1. CAMSAP'05, 2005, 129-132.

[3] L. Qi, Eigenvalues of a real supersymmetric tensor, *J. Symbolic Comput.* 40 (2005) 1302-1324.

[4] L. Qi,  $H^+$ -eigenvalues of Laplacian and signless Laplacian tensors, *Commun. Math. Sci.* 12 (6) (2014) 1045-1064.

[5] L. Liu, L. Kang, S. Bai, Bounds on the spectral radius of uniform hypergraphs, *Discrete Applied Mathematics* 259. 10.1016/j.dam.2018.12.007.

[6] S. Friedland, S. Gaubert, L. Han, Perron-Frobenius theorem for nonnegative multilinear forms and extensions, *Linear Algebra Appl.* 438 (2013) 738-749.

[7] Y. N. Yang, Q. Z. Yang, Further Results for Perron-Frobenius theorem for nonnegative tensors, *SIAM Journal on Matrix Analysis and Applications* 31 (2010) 2517-2530.

[8] J. Shao, A general product of tensors with applications, *Linear Algebra Appl.* 439 (2013) 2350-2366.

[9] G. Yi, *Tensor spectral properties of uniform hypergraphs*, Fuzhou University 2014.

[10] Y. X. Yuan, M. Zhang, M. Lu, Some upper bounds on the eigenvalues of uniform hypergraphs, *Linear Algebra Appl.* 2015.

[11] L. Qi, Z. Luo, *TENSOR ANALYSIS*, Society for Industrial and Applied Mathematics Philadelphia, 2017.

[12] S. R. Bulò, M. Pelillo, *New bounds on the clique number of graphs based on spectral hypergraph theory*, Springer, Berlin, Heidelberg, 2009.

[13] Y. Yang, Q. Yang, Further results for Perron-Frobenius theorem for nonnegative tensors, *SIAM J. Matrix Anal. Appl.* 31 (5) (2010) 2517-2530.

[14] Y. Yang, Q. Yang, On some properties of nonnegative weakly irreducible tensors, arXiv: 1111.0713v3, 2011.

[15] V. T. Sós, E. G. Straus, Extremals of functions on graphs with applications to graphs and hypergraphs, *J. Combin. Theory Ser. B*, 32 (1982) 246-257.