

Biofilm Formation Between *Bacillus Subtilis* and *Escherichia Coli* K-12 Strains at Acidic and Oxidative Stress

Ivo Ganchev

Department of General Microbiology, The Stephan Angeloff Institute of Microbiology, Bulgarian Academy of Sciences, Sofia, Bulgaria

Email address:

ivotg@abv.bg

To cite this article:

Ivo Ganchev. Biofilm Formation Between *Bacillus Subtilis* and *Escherichia Coli* K-12 Strains at Acidic and Oxidative Stress. *Science Journal of Chemistry*. Vol. 7, No. 1, 2019, pp. 15-18. doi: 10.11648/j.sjc.20190701.13

Received: September 30, 2018; Accepted: October 24, 2018; Published: April 18, 2019

Abstract: Biofilms constitute the predominant microbial style of life in natural and engineered ecosystems. At the harsh environmental conditions microorganisms accumulate reactive oxygen species (ROS), potentially encountering a dangerous condition called oxidative stress. An investigation into the mechanisms activated by biofilms in response to different oxidative stress levels could have important consequences from ecological and economic points of view, and could be exploited to propose alternative strategies to control microbial virulence and deterioration. In this respect, the aim of this study is to evaluate the influence of temperature and pH of the medium, its osmolality and the presence of heavy metal ions on the growth of the biomass of biofilm formed by *B. subtilis* 170 and *E. coli* K-12 strains and their relationship in them. This study used the methods for a static cultivation of biofilm by co-cultures of *B. subtilis* 170 and *E. coli* K-12 strains and determination of colony forming units in their structures to evaluate the relationships between them. The results of the present study show that temperature of 20°C and pH-value in the range of 5.0-6.0 help to maximum growth of biomass of biofilm formed by co-cultivation of *B. subtilis* 170 and *E. coli* 1655 strains. The competitive relationships are observed at a temperature of 20°C, at pH-value in the range from 5.0 to 6.0 at low osmolality of the medium of 100 to 150 mM, at content of Fe^{+2} from 5 μM to 100 μM . The increase of the temperature above 30°C and the pH of the medium, the high osmolality of 200 mM triggers antagonistic interactions between *B. subtilis* 170 and *E. coli* K-12 strains, while at content of Fe^{+2} of 50 μM was observed symbiosis in the structure of biofilms in this study.

Keywords: *B. Subtilis* 170, *E. Coli* 1655, Biofilm, Competition, Antagonism

1. Introduction

Bacteria were distributed in nature in the form of biofilms that are formed with the participation of two or more species [1-3]. The basis of their formation and their resistance toward various effects of the environment in the nature stand interactions between microbial species that include competition for nutrients of the composition of the environment, symbiosis and antagonism [2]. Therefore, the structure of biofilms and the composition of the microbial community in them are determined by the fluctuations in values of environmental factors [2-4].

Harsh environmental conditions such as high temperature, osmolality and pH of the medium, presence of heavy metal ions (Fe^{2+} or Fe^{3+} or Cu^{+}) creates conditions for formation

and accumulation of intracellular free radicals such as superoxide anion, hydrogen peroxide, singlet oxygen, hydroxyl radicals (oxidative stress), which in the high (lethal) concentrations have a cytotoxic effect and prevent biofilm formation process. Main mechanisms for protection of microbial species against their impact during the formation of biofilms is concluded in the presence of complex genetically regulatory network, responsible for the transition of cells from plankton state to attached mode of existence, and the formation of extracellular matrix, whose content is dominated by extracellular polysaccharides [5].

The aim of this study is to evaluate the influence of temperature and pH of the medium, its osmolality and the presence of heavy metal ions on the growth of the biomass of biofilm formed by *B. subtilis* 170 and *E. coli* K-12 strains

and their relationship in them.

2. Material and Methods

2.1. Bacterial Strains

In this study were used *B. subtilis* 170 from collection, deposited in National Bank of Industrial Microorganisms and Cell Cultures, Sofia, Bulgaria, and *E. coli* 1655 from the collection of the Institute of Molecular Biotechnology, Jena, Germany. All strains were inoculated into 9 ml of a liquid culture medium and incubated at 37°C for a period of 18 hours before the beginning of each determination.

2.2. Method of Static Cultivating of Biofilms

Liquid cultures were diluted in 5 ml medium M63 in different ratio between KH_2PO_4 and K_2HPO_4 , so that the pH of the medium to be 5.0; 6.0; 7.0; 8.0. Study of influence of the temperature was carried out by incubating the cultures at 20°C, 30°C, 35°C and 40°C in medium M63 (0.02 M KH_2PO_4 , 0.04 M K_2HPO_4 , 0.02 M $(\text{NH}_4)_2\text{SO}_4$, 0.1 mM MgSO_4 and 0.04 M glucose).

Immediately after that was carried cultivation of biofilms in environments M63, containing NaCl of 100 mM, 150 mM and 200 mM for taking into account the effects of osmolality on their growth. This was followed by carrying out a series of cultivation of biofilms in modified M63 medium to which is add $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, so that the content of Fe^{+2} to be 5 μM , 50 μM and 100 μM .

2.3. Determination of Biomass Growth of Biofilms

After incubation at 20°C for 24 h from each well separated carefully culture medium, after washing three times with saline solution staining of biofilms in half of the wells was carried out with 200 μl 0,1% solution of crystal violet over a period of 15 min, gently release and washing of dye in each well, then were added 200 μl of 70% ethanol. Immediately after that it was determined the optical density at 540 nm on "ELLIZA Reader ELTA 90".

2.4. Determining of Number of Colonies in the Biofilms

In the second half of the wells were added 200 μl 0.85%

sodium chloride solution. Using a sterile knife the biofilm was separated carefully. For each variant were made in 6 wells. Solutions with a release biofilm were collected in sterile eppendorff tubes. The number of colonies in the biofilms was determined according to the method of Koch [6] after inoculation on Meat Peptone Agar (NCIPD-Bull Bio, Sofia, Bulgaria) to evaluate of total number of colonies and on MacConkey agar (NCIPD-Bull Bio, Sofia, Bulgaria) for counting of number of colonies of *E. coli* 1655 strain.

2.5. Statistical Treatment of Results

All experiments were performed three times, presented results represent the average of three independent each other determinations. It was used a Student test, the differences of individual values were considered statistically significant at $p < 0.05$.

3. Results and Discussion

Biofilms, which are formed by the participation of different bacterial species are highly distributed in nature. In the process of their forming establish different relationships between microorganisms that include competition for nutrients of the composition of the medium, antagonism or symbiosis. The process of formation of these organized structures on different surfaces strongly conditioned on the fluctuations in environmental factors [1].

The strains of the *B. subtilis* and *E. coli* species during the process of independent and co-culturing of biofilms maintain intracellular pH in narrow range from pH 7,2 to pH 7,8 at changing the pH of culture medium from 5.0 to 8,0 (Figure 1). The pH - value of the medium in the range of from pH 5,0 to pH 6,0 provides the presence of cells of *E. coli* K-121655 strain in the structure of the biofilm in association with *B. subtilis* 170 strains, which may be associated with the presence of potassium ions that contribute to the sustainability of cells against acid stress. However, the percentage of their population in the resulting biofilms is relatively small compared to the number of colonies of *B. subtilis* 170 strain, which indicates a competitive relationship (Table 1).

Table 1. Influence of the pH and temperature on the number of colonies of *B. subtilis* 170 and *E. coli* K-12 1655 strains in the structure of monospecies and mixed biofilms.

№	pH of medium	Number in biofilms of <i>B. subtilis</i> 170 cfu/ml	Number in biofilms of <i>E. coli</i> K-12 1655, cfu/ml	Number in mixed biofilms cfu/ml	
				<i>B. subtilis</i> 170	<i>E. coli</i> K-12 1655
1.	5,0	9,05±0,18.10 ⁶	1,62±0,12.10 ⁶	9,55±0,13.10 ⁶	3,16±0,12.10 ⁶
2.	6,0	8,11±0,07.10 ⁶	0,26±0,03.10 ⁶	35,91±0,05.10 ⁶	0,40±0,05.10 ⁶
3.	7,0	0,56±0,07.10 ⁶	0	10,95±0,13.10 ⁶	0
4.	8,0	0,23±0,03.10 ⁶	0	6,18±0,20.10 ⁶	0

№	Temperature, °C	Number in biofilms of <i>B. subtilis</i> 170 cfu/ml	Number in biofilms of <i>E. coli</i> K-12 1655, cfu/ml	Number in mixed biofilms cfu/ml	
				<i>B. subtilis</i> 170	<i>E. coli</i> K-12 1655
1.	20	6.65±0.1.10 ⁶	0,4±0.0.10 ⁶	46.31±0.33.10 ⁶	6.01±0.03.10 ⁶
2.	30	7.06±0.07.10 ⁶	0	32,4±0.13.10 ⁶	0
3.	35	0.63±0.05.10 ⁶	0	4.65±0.05.10 ⁶	0
4.	40	0.56±0.08.10 ⁶	0	0.42±0.07.10 ⁶	0




Figure 1. Effect of pH and temperature on the growth of biofilms: ■ *B. subtilis* 170; ■ *B. subtilis* 168; ■ *E. coli* 1655; ■ *B. subtilis* 170+ *E. coli* 1655; □ *B. subtilis* 168+ *E. coli* 1655.

Higher pH - value of culture media are created conditions for the emergence of antagonistic relationships, as result the biofilms are formed only with the participation of cells of the *B. subtilis* strains. The obtained values for the structure of the microbial population in the biofilms formed at a pH from 7,0 to 8,0 due to the fact that maintenance of pH - homeostasis in *E. coli* K-12 1655 strains in alkaline media is due to low activity of the system of antiport between sodium ions and protons, which determines their weak adaptive capacity under conditions of cultivation. On the another hand, the antagonistic activity of the *B. subtilis* 170 strain is associated with the creation of optimal conditions for action of secreted antimicrobial proteins whose biosynthesis is a function of the temperature of cultivation.

The temperature of cultivation influences on the growth of the biomass of biofilms formed by the participation of strains of *B. subtilis* and *E. coli* species, highest values are reported at 20°C (Figure 1). The reduction of the total biomass by increasing of temperature in the range from 30°C to 40°C is probably associated with the process of oxidation processes that leads to damage of the cell and cell proteins, chain of DNA [5-8]. Moreover, in the structure of the biofilm is established competitive relationships between the *B. subtilis* 170 and *E. coli* K-12 1655 strains at a temperature of cultivation of 20°C, while the increasing of its value in the range from 20 to 40°C creates conditions for the development of antagonistic relationships, as a result of which the structure of the biofilm is dominated by cells of *B.*

subtilis 170 strain, but the number of their colonies is decreasing from $32,4 \pm 0.13 \cdot 10^6$ cfu/ml to $0.42 \pm 0.07 \cdot 10^6$ cfu/ml in the structure of the biofilm formed by co-cultivation of pair of *B. subtilis* 170 and *E. coli* K-12 1655 strains (Table 1).

The osmolality of the medium influences on biofilm formation of a large number of bacterial species [9]. The increase of content of sodium chloride from 100 mM to 200 mM results in a reduction of growth of the biomass of biofilms of *B. subtilis* 170 strain (Figure 2). The significant increase of numbers of cell population in the biofilm structure of *E. coli* K-12 1655 strain from $1 \pm 0,2 \cdot 10^3$ cfu/ml to $12,53 \pm 0,21 \cdot 10^3$ cfu/ml probably relates to the activation of transcription of the *pga* operon, which is responsible for the biosynthesis of proteins, necessary for the formation of polyacetylglucosamine, that is the part of the composition of the matrix [9,10]. Joint development with *B. subtilis* 170 strain doesn't determine its dominate role in the formation of structures of biofilms, given to the repression of genes for curly-fringe at the influence of products of life activity of strains of *B. subtilis* species. Accordingly, the content of sodium chloride of 200 mM in the medium leads to absent of cells of the *E. coli* K-12 1655 strain in biofilms at his association with the strains of *B. subtilis* species, which is an indication to perform the transition from competitive relationships to antagonism at increasing osmolality of the medium (Table 2).




Figure 2. Effect of osmolality and Fe^{+2} of the medium on the growth of biofilms: ■ *B. subtilis* 170; ■ *B. subtilis* 168; ■ *E. coli* 1655; ■ *B. subtilis* 170+ *E. coli* 1655; □ *B. subtilis* 168+ *E. coli* 1655.

Table 2. Effect of concentration of sodium chloride in the medium on number of colonies of *B. subtilis* 170 and *E. coli* K-12 1655 strains in the structure of monospecies and mixed biofilms.

№	Concentration of NaCl, mM	Number in biofilms of <i>B. subtilis</i> 170 cfu/ml	Number in biofilms of <i>E. coli</i> K-12 1655, cfu/ml	Number in mixed biofilms cfu/ml	
				<i>B. subtilis</i> 168	<i>E. coli</i> K-12 1655
1.	100	36,5±0,62.10 ³	1±0,2.10 ³	116,43±0,97.10 ³	31,56±0,32.10 ³
2.	150	10,26±.10 ³	0,22±.10 ³	39,46±0,72.10 ³	21,13±0,21.10 ³
3.	200	10,3±0,26.10 ³	12,53±0,21.10 ³	13,5±0,3.10 ³	0
№	Concentration of Fe ²⁺ , µM	Number in biofilms of <i>B. subtilis</i> 170 cfu/ml	Number in biofilms of <i>E. coli</i> K-12 1655, cfu/ml	Number in mixed biofilms cfu/ml	
				<i>B. subtilis</i> 170	<i>E. coli</i> K-12 1655
1.	5	4.63±0.3.10 ³	4.83±0.30.10 ³	42.96±0.40.10 ³	5.23±0.15.10 ³
2.	50	56.23±0.21.10 ³	19.46±0.21.10 ³	53.03±0.49.10 ³	40.26±0.25.10 ³
3.	100	4.5±0.2.10 ³	5.43±0.32.10 ³	102.93±0.29.10 ³	8.86±0.15.10 ³

On the other hand, iron belongs to the trace elements, which content influences on the nature of relationships between strains of *B. subtilis* and *E. coli* species during formation of biofilms [11]. The change of content of ferrous ions in the medium in the range from 5 µM to 100 µM is accompanied by the increase in biomass of mixed biofilms (Figure 2), optical density at 540 nm changed in a range from 0,163 to 0,184 as a result of the interaction between *B. subtilis* 170 and *E. coli* K-12 1655 strains. Low concentration of ferrous ions than 5 µM in the medium catalyzes processes of intracellular accumulation of •OH, which reacts with biomakromolecules of cells [12] and provides a connection to cell surface protein in *E. coli* K-12 1655 strains and repression of genes for colonic acid biosynthesis, which explains their poor spread in the structure of biofilms at their interaction with *B. subtilis* 170 strain [10] (Table 2). The value of ferrous ion of 50 µM stimulated symbiosis between investigated pair strains which may be attributed to the establishment of a balance between the generation of free radicals in cells of the both strains and the activity of superoxide dismutase enzyme systems, resulting in lowering the cytotoxic effect of the generated free radicals at both strains [13-15].

4. Conclusions

The effect of temperature of cultivation, pH, osmolality, presence of Fe²⁺ on the growth of biofilms formed by a mixed population of strains of *B. subtilis* and *E. coli* species was investigated. Temperature of 20°C and the pH value of the medium in the range from pH 5,0 to pH 6,0 maximally stimulate growth of biofilm biomass of *B. subtilis* 170 and *E. coli* K-12 1655 strains.

Changes of pH-value, the temperature of cultivation, osmolality of medium and the presence of iron ions affect the nature of the interaction between strains of *B. subtilis* and *E. coli* species in presen study, competitive relationships are observed at a temperature of 20°C, the pH in the range from 5.0 to 6.0 at low osmolality of the medium of 100 to 150 mM, at content of Fe²⁺ from 5 µM to 100 µM. The increase of the temperature above 30°C and pH of the medium, high osmolality of 200 mM triggers antagonistic interactions between test pairs strains, while at content of Fe²⁺ of 50 µM

was observed symbiosis during process of formation of biofilms.

References

- [1] Elias S., E. Banin, FEMS Microbiol Rev, 36 (5), 990–1004 (2012).
- [2] Moons P, Chris W. Michiels, A. Aertsen, Critical Reviews in Microbiology, 35(3), 157–168 (2009).
- [3] Huang R., M. Li, R. Gregory, Virulence, 2 (5), 435-444 (2011).
- [4] Martinez, R. Kitko, J. Patrick Mershon, H.. Adcox, K.. Malek, M. Berkmen, J. L. Slonczewski, Applied and Environmental Microbiology, 78(10), 3706–3714(2012).
- [5] Gambino M., F. Cappitelli, Biofouling, 32(2), 167–178 (2016).
- [6] Grudeva V., P. Moncheva, S. Naumova, B. Gocheva, T. Nedeva, S. Antonova-Nikolova (2006). In: Manual of Microbiology; ("St. Kliment Ohridski" University Edition), Sofia, 146-153.
- [7] Murata M, Fujimoto H, Nishimura K, Charoensuk K, Nagamitsu H, Raina S, Kosaka T, Oshima T, Ogasawara N, Yamada M., PloS One., 6, 20-63 (2011).
- [8] Chen J, Shen J, Solem C, Jense, Appl Environ Microbiol., 79, 6140–6147(2013).
- [9] Goller, C., X. Wang, Y. Itoh, T. Romeo, J. Bacteriol. 188, 8022–8032 (2006).
- [10] Karatan E., P. Watnick, Microbiology and Molecular Biology Reviews, 73 (2), 310–347 (2009).
- [11] Ryder C., M. Byrd, and D. J. Wozniak, Curr. Opin. Microbiol., 10,644–64 (2007).
- [12] Bokare AD, Choi W., J Hazard Mat, 275, 121–135(2014).
- [13] Cabisco E, Tamarit J, Ros J., Intern Microbiol., 3, 3–8(2000).
- [14] Green J, Paget MS., Nat Rev Microbiol., 2, 954–966 (2004).
- [15] Imlay J., Curr Opin Microbiol., 24, 124–131 (2015).